



## Secondhand tobacco smoke concentrations in motor vehicles: a pilot study

M R Jones, A Navas-Acien, J Yuan and P N Breyse

*Tob. Control* 2009;18;399-404; originally published online 25 Aug 2009;  
doi:10.1136/tc.2009.029942

---

Updated information and services can be found at:  
<http://tobaccocontrol.bmj.com/cgi/content/full/18/5/399>

---

*These include:*

### References

This article cites 28 articles, 14 of which can be accessed free at:  
<http://tobaccocontrol.bmj.com/cgi/content/full/18/5/399#BIBL>

### Rapid responses

You can respond to this article at:  
<http://tobaccocontrol.bmj.com/cgi/eletter-submit/18/5/399>

### Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

---

### Notes

---

To order reprints of this article go to:  
<http://journals.bmj.com/cgi/reprintform>

To subscribe to *Tobacco Control* go to:  
<http://journals.bmj.com/subscriptions/>

# Secondhand tobacco smoke concentrations in motor vehicles: a pilot study

M R Jones,<sup>1,2</sup> A Navas-Acien,<sup>1,2,3</sup> J Yuan,<sup>2</sup> P N Breyse<sup>2,3</sup>

<sup>1</sup> Department of Epidemiology, Johns Hopkins University Bloomberg School of Public Health, Baltimore, Maryland, USA; <sup>2</sup> Department of Environmental Health Sciences, Johns Hopkins University Bloomberg School of Public Health, Baltimore, Maryland, USA; <sup>3</sup> Institute for Global Tobacco Control at the Johns Hopkins University Bloomberg School of Public Health, Baltimore, Maryland, USA

Correspondence to:  
Dr Ana Navas-Acien,  
Department of Environmental  
Health Sciences, Johns Hopkins  
Bloomberg School of Public  
Health, 615 N Wolfe Street,  
Office W7033B, Baltimore, MD  
21205, USA; [anavas@jhsph.edu](mailto:anavas@jhsph.edu)

Received 5 February 2009  
Accepted 30 June 2009

## ABSTRACT

**Context:** Motor vehicles represent important micro-environments for exposure to secondhand smoke (SHS). While some countries and cities have banned smoking in cars with children present, more data are needed to develop the evidence base on SHS exposure levels in motor vehicles to inform policy and education practices aimed at supporting smoke-free motor vehicles when passengers are present.

**Objective:** To assess exposure to secondhand tobacco smoke in motor vehicles using passive airborne nicotine samplers.

**Methods:** 17 smokers and five non-smokers who commute to and from work in their own vehicle participated. Two passive airborne nicotine samplers were placed in each vehicle for a 24-hour period, one at the front passenger seat headrest and the other in the back seat behind the driver. At the end of the sampling period, airborne nicotine was analysed by gas chromatography.

**Results:** Median (IQR) air nicotine concentrations in smokers' vehicles were 9.6 µg/m<sup>3</sup> (5.3–25.5) compared to non-detectable concentrations in non-smokers' vehicles. After adjustment for vehicle size, window opening, air conditioning and sampling time, there was a 1.96-fold increase (95% CI 1.43 to 2.67) in air nicotine concentrations per cigarette smoked.

**Conclusions:** Air nicotine concentrations in motor vehicles were much higher than air nicotine concentrations generally measured in public or private indoor places, and even higher than concentrations measured in restaurants and bars. These high levels of exposure to SHS support the need for education measures and legislation that regulate smoking in motor vehicles when passengers, especially children, are present.

Involuntary exposure to secondhand smoke accounts for thousands of cases of respiratory, cardiovascular and cancer deaths in the United States each year.<sup>1</sup> To prevent mortality and morbidity caused by secondhand smoke, smoke-free regulations banning smoking in public places are being passed in cities and countries around the world.<sup>2–5</sup> Public education is also being implemented to discourage smoking inside the home.<sup>4</sup> Indeed, households enforcing smoke-free home rules in the US increased from 43% in 1992–3 to 72% in 2003.<sup>5</sup> With commuting lengths increasing and the importance of car pooling, it is vital to consider the health risks of exposure to secondhand smoke to motor vehicle passengers who travel with a smoker.

While scarce data quantifying secondhand smoke exposure levels in cars exist,<sup>6–9</sup> motor vehicles represent important microenvironments for secondhand smoke exposure—first, because there is no risk-free level of secondhand smoke

exposure<sup>1</sup>; second, because owing to the confined space within a motor vehicle, tobacco smoke concentrations increase rapidly<sup>10</sup> posing serious health risks for passengers who are repeatedly exposed. Epidemiological and mechanistic studies have shown that short-term secondhand smoke exposure can have a substantial adverse impact on the cardiovascular system, similar to active smoking.<sup>11</sup> Peak secondhand smoke exposure can also impact lung function decline in asthmatic children.<sup>12</sup> Regulatory measures banning smoking in motor vehicles, however, have only been passed in a small number of countries, states and municipalities, and are limited to the protection of children of varying ages.<sup>6–13</sup>

This pilot study measured air nicotine concentrations in motor vehicles of smokers and non-smokers while they commute to and from work. The application of the protocol under actual driving conditions rather than experimental conditions, as was done in previous studies,<sup>7–9</sup> may better reflect actual exposure concentrations during driving. Given available evidence on the lack of protection of ventilation measures in indoor places,<sup>14</sup> we hypothesised that ventilation measures in the vehicle, such as having windows down or air conditioning on, may be ineffective in providing complete protection from exposure to secondhand smoke to non-smoking passengers.

## METHODS

### Design and population

This exposure survey used air nicotine passive samplers to characterise passenger exposure to secondhand smoke in motor vehicles. During the summer of 2008, employees of a large medical institution in Baltimore, Maryland, who congregated outdoors around their workplace were invited to participate. To be eligible, the employee had to regularly operate a vehicle and have a daily commute that lasted 30 minutes or more. Smokers also had to regularly smoke within the vehicle. In total, 29 drivers were recruited, 24 smokers and five non-smokers. Each driver received two air nicotine samplers. A total of 17 smokers (71%) and five non-smokers (100%) returned the nicotine samplers, leaving 44 air nicotine samplers in the present study. Reasons for not returning the samplers after initial recruitment included health problems (n = 2), lost samplers (n = 1) and unknown reasons (n = 4). Smokers initially recruited and smokers returning samplers were similar for all participant and vehicle characteristics evaluated at recruitment (table 1). Participation in the study was voluntary. Written and oral consent was obtained from all participants. The study protocol and consent forms were

**Table 1** Summary of participant characteristics

|   | Smokers recruited | Smokers returning monitors | Non-smokers recruited and returning monitors |
|---|-------------------|----------------------------|--|
| Number  | 24                | 17                         | 5  |
| Age, mean (SD)  | 39 (11)           | 38 (12)                    | 37 (11)                                      |
| Sex, % male   | 33                | 47                         | 60   |
| Education (%)   |                   |                            |  |
| High school (<13 years)   | 26                | 31                         | 20   |
| College + (≥13 years)   | 74                | 69                         | 80   |
| Vehicle activity  |                   |                            |  |
| Days/week driving vehicle, mean (SD)                            | 7 (1)             | 7 (1)                      | 5 (2)  |
| Hours/day driving vehicle, mean (SD)                            | 2 (1)             | 2 (1)                      | 1 (1)  |
| Active smoking (%)  |                   |                            |  |
| Current smoker  | 100               | 100                        | 0  |
| Smokes daily  | 96                | 94                         | –  |
| Smokes less than daily  | 4                 | 6                          | –  |
| Smoking in vehicle (%)  |                   |                            |  |
| Every day   | 83                | 76                         | –  |
| Some days   | 17                | 24                         | –  |
| Do not smoke in the vehicle                                     | –                 | –                          | 100  |
| Vehicle size* (%)   |                   |                            |  |
| Compact/small   | 42                | 35                         | 60   |
| Mid-size/large  | 58                | 65                         | 40   |
| Number of passengers usually occupying the vehicle†             |                   |                            |  |
| 0   | 58                | 58                         | 40   |
| 1   | 17                | 12                         | 40   |
| ≥2  | 25                | 30                         | 20   |
| Number of children usually occupying the vehicle                |                   |                            |  |
| 0   | 71                | 71                         | 60   |
| ≥1  | 29                | 29                         | 40   |
| Have rules in car that restrict smoking (%)                     | 79                | 82                         | 100  |
| Smoking rules/restrictions (%)                                  |                   |                            |  |
| Not allowed in vehicle  | 0                 | 0                          | 100  |
| Allowed only with use of ventilation                            | 20                | 21                         | –  |
| Allowed only without passengers                                 | 15                | 7                          | –  |
| Allowed with ventilation and no passengers                      | 65                | 71                         | –  |
| Driver opinion (%)  |                   |                            |  |
| Smoking poses a health risk to passengers                       | 100               | 100                        | 100  |
| Being unable to smoke in the car would help you to quit smoking | 53                | 53                         | –  |
| Vehicles should be smoke-free on a voluntary basis              | 93                | 93                         | 60   |
| Vehicles should be smoke-free by regulation                     | 7                 | 7                          | 60   |

\*Vehicle types included compact and midsize cars, minivans, and sport utility vehicles.

†Includes both children and adult passengers combined.

approved by the Johns Hopkins Bloomberg School of Public Health institutional review board.

### Nicotine monitoring

Air nicotine concentrations in each vehicle were estimated by passive sampling of vapour-phase nicotine using a sampler originally developed by Hammond and Leaderer.<sup>16</sup> The samplers were assembled from a 37-mm sampling cassette containing a filter treated with sodium bisulfate and covered by a polycarbonate diffusion membrane at the Secondhand Smoke Exposure Assessment Laboratory of the Johns Hopkins Institute for Global Tobacco Control.

Two nicotine samplers were placed in each vehicle for a 24-hour period in a sealed container. To evaluate levels of secondhand smoke exposure expected for actual passengers, one sampler was placed at the base of the front passenger seat headrest and the other was placed on the lateral hanger in the back seat behind the driver. Drivers were instructed to open the samplers during travelling time within the 24-hour sampling period and to keep them closed otherwise. All participants

received and were instructed to use latex gloves when handling the air samplers. At the end of the sampling period, the collected nicotine was extracted from the filters and analysed by gas chromatography using a nitrogen phosphorous detector.

The time-weighted concentration of airborne nicotine was calculated by dividing the amount of nicotine collected by the filter ( $\mu\text{g}$ ) per volume of air sampled ( $\text{m}^3$ ). Volume sampled was calculated by multiplying the time sampled (provided by the participant) by the effective sampling rate of the sampler ( $25 \text{ ml/min}$ ).<sup>16</sup> We assumed that the sampling rate was not affected by changes in face velocity caused by open windows. Blanks were used to determine the blank-corrected nicotine concentrations and to calculate the nicotine limit of detection ( $0.0026 \mu\text{g/ml}$  corresponding to  $0.22 \mu\text{g/m}^3$  for 1 hour of sampling time). For samples below the limit of detection a value half the limit of detection was assigned.

### Data collection

At recruitment, participants were asked to respond to a short questionnaire requesting information on age, sex, education,

smoking behaviour and vehicle characteristics (vehicle make, model, year and length of ownership) and occupancy (adult, children and smoking passengers). The vehicle types represented included compact and midsize cars as well as minivans and sport utility vehicles (SUVs). For analysis, the vehicles were grouped into size categories of compact/small and midsize/large vehicles. At the end of the sampling period, each driver was requested to report information on time spent in the vehicle, ventilation use, number of cigarettes smoked and vehicle occupancy for any trips occurring during the 24-hour study. Drivers were also asked about knowledge and attitudes regarding the health risks of secondhand smoke, secondhand smoke regulations and the potential impact of motor vehicle smoking bans on their ability to quit smoking.

### Statistical analyses

Descriptive analyses were stratified by participant smoking status. Air nicotine concentrations from samplers placed on the front passenger seat headrest (geometric mean 3.9, 95% confidence interval (CI) 1.5 to 10.1) and back seat behind the driver (geometric mean 2.8, 95% CI 1.1 to 7.3) were comparable ( $p = 0.12$ ) and concentrations are presented as the average of the two air nicotine samplers in each vehicle. Air nicotine concentrations were described using the median and interquartile range (IQR) overall and by participant and motor vehicle characteristics. To compare nicotine concentrations by number of cigarettes smoked within the vehicle, vehicle size and other travelling conditions, we computed crude and multivariable adjusted ratios of the geometric mean of nicotine concentrations and its 95% CI using linear regression models on log-transformed nicotine. The crude and adjusted ratios (95% confidence interval) of geometric means by motor vehicle characteristics were estimated by exponentiating the coefficients and standard errors from the linear regression models on log-transformed nicotine. Analyses were conducted using Stata version 9.0.

## RESULTS

### Participant and motor vehicle characteristics


Mean age of study participants who returned the nicotine air samplers ( $n = 22$ ) was 37 (SD 12) years, 50% were men and 71% had college education or more, with little differences by smoking status (table 1). On average, smoking participants drove 7 (SD 1) days a week and 2 (SD 1) hours/day compared to 5 (SD 2) days a week and 1 (SD 1) hour/day for non-smoking drivers. Smoking drivers smoked in their vehicle on a daily basis and all smoked filtered cigarettes. The number of cigarettes smoked during the sampling period (mean 4.65, SD 2.34) was similar to the average number reported as being smoked daily in the vehicle (mean 5.09, SD 2.75). The size of the vehicle was compact/small for 35% and 60% of smoking and non-smoking drivers, respectively. In 45% of motor vehicles there was at least one passenger in addition to the driver and in 31% there was at least one child passenger, with no major differences by smoking status. None of the non-smoking drivers permitted smoking in the vehicle. All smoking drivers allowed smoking within the vehicle, although 71% of them allowed it only without passengers and with use of ventilation.

### Air nicotine concentrations

The median (IQR) sampling time during the 24-hour period was 105 (80–135) minutes. Thirteen samplers had air nicotine concentrations below the limit of detection, including 10

samplers from five non-smokers and three samplers from two smokers. The smoking driver with both front and rear samplers below the limit of detection reported smoking a total of three cigarettes and having the windows down more than halfway during the overall 40-minute sampling time (the shortest commute of all of the drivers in the study). For the other smoking driver, only the rear sampler was below the limit of detection although the nicotine concentration in the front sampler was also relatively low ( $3.2 \mu\text{g}/\text{m}^3$ ). The vehicle was a large SUV with ventilation on and a total of five cigarettes were smoked during 110 minutes of sampling time.

Median (IQR) air nicotine concentrations in smokers' vehicles were  $9.6 \mu\text{g}/\text{m}^3$  (5.3–25.5) compared to undetectable in non-smokers' vehicles during the sampling period (table 2). The maximum nicotine concentration,  $128.1 \mu\text{g}/\text{m}^3$ , occurred in a small vehicle with a total of eight cigarettes smoked during the 105-minute sampling period. The driver reported having the windows closed and used air conditioning throughout the sampling period. Air nicotine concentrations increased with increasing number of cigarettes smoked during the sampling period (median air nicotine  $8.3 \mu\text{g}/\text{m}^3$  and  $12.5 \mu\text{g}/\text{m}^3$  for one to three cigarettes and four or more cigarettes smoked, respectively) (fig 1). Air nicotine concentrations were 1.96 times higher (95% CI 1.43 to 2.67) per cigarette smoked after



**Figure 1** Air nicotine concentrations\* ( $\mu\text{g}/\text{m}^3$ ) in motor vehicles by number of cigarettes smoked. \*Average of front and rear air nicotine concentrations. (A) Box plots where horizontal lines within boxes represent medians; boxes, interquartile ranges; bars, values within 1.5 times the interquartile range; solid circles, outlying data points. (B) Scatterplot where the line represents the linear dose-response relation between the number of cigarettes smoked and the  $\log_{10}$  nicotine concentrations;  $r$  is the Spearman correlation coefficient.

## Research paper

**Table 2** Air nicotine concentrations\* ( $\mu\text{g}/\text{m}^3$ ) by vehicle and travelling characteristics

|                               | No | Median (IQR)<br>smokers | No | Median (IQR)<br>non-smokers† | Crude ratio (95% CI) | Adjusted‡ ratio (95% CI) |
|-------------------------------|----|-------------------------|----|------------------------------|----------------------|--------------------------|
| Overall                       | 17 | 9.6 (5.3–25.5)          | 5  | $\leq 0.1$                   | –                    | –                        |
| No of cigarettes smoked       |    |                         |    |                              |                      |                          |
| 0                             | 0  | –                       | 5  | $\leq 0.1$                   | 1.00 (reference)     | 1.00 (reference)         |
| 1–3                           | 4  | 8.3 (3.7–10.2)          | 0  | –                            | 35.75 (7.83 to 163)  | 72.40 (15.8 to 332)      |
| $\geq 4$                      | 13 | 12.5 (5.3–31.7)         | 0  | –                            | 122.28 (37.2 to 403) | 293.56 (87.2 to 981)     |
| Per cigarette smoked          |    | –                       |    | –                            | 1.79 (1.41 to 2.27)  | 1.96 (1.43 to 2.67)      |
| Size of car¶                  |    |                         |    |                              |                      |                          |
| Midsize/large                 | 11 | 7.5 (5.3–12.5)          | 2  | $\leq 0.1$                   | 1.00 (reference)     | 1.00 (reference)         |
| Compact/small                 | 6  | 32.3 (7.0–43.9)         | 3  | $\leq 0.1$                   | 0.76 (0.11 to 5.51)  | 4.34 (1.44 to 13.0)      |
| Windows opened§               |    |                         |    |                              |                      |                          |
| Less than halfway             | 7  | 9.6 (5.3–12.5)          | 4  | $\leq 0.1$                   | 1.00 (reference)     | 1.0 (reference)          |
| More than halfway             | 10 | 13.0 (5.3–31.7)         | 1  | $\leq 0.1$                   | 2.94 (0.44 to 19.5)  | 0.40 (0.13 to 1.21)      |
| Air conditioning              |    |                         |    |                              |                      |                          |
| Off                           | 8  | 7.4 (6.1–14.0)          | 1  | $\leq 0.1$                   | 1.00 (reference)     | 1.0 (reference)          |
| On                            | 9  | 12.5 (5.3–39.1)         | 4  | $\leq 0.1$                   | 0.81 (0.11 to 5.84)  | 0.44 (0.13 to 1.50)      |
| Total sampling time (minutes) |    |                         |    |                              |                      |                          |
| $\leq 60$                     | 3  | 7.0 (0.3–7.3)           | 0  | –                            | 1.00 (reference)     | 1.0 (reference)          |
| $>60$                         | 14 | 11.7 (5.3–31.7)         | 5  | $\leq 0.1$                   | 1.48 (0.09 to 25.3)  | 6.88 (1.32 to 35.9)      |

\*Average of front and rear air nicotine concentrations.

†Air nicotine concentrations of all five vehicles of non-smokers were below the limit of detection.

‡Ratios adjusted for all other variables with adjustment for cigarette smoked conducted using the categorical variable, except for the model per cigarette smoked.

¶Vehicle types included compact and midsize cars, minivans and SUVs.

§For most of the trip.

adjustment for vehicle size, window opening, air conditioning and length of sampling time with small differences before adjustment (ratio 1.79; 95% CI 1.41 to 2.27) (table 2).

Among smoking drivers, median air nicotine concentrations were higher in compact/small vehicles (median  $32.3 \mu\text{g}/\text{m}^3$ ) compared to midsize/large vehicles (median  $7.5 \mu\text{g}/\text{m}^3$ ) (table 2). After adjustment for number of cigarettes and other driving condition characteristics, air nicotine concentrations in smokers' vehicles were 4.34-fold time increased in compact/small vehicles compared to midsize/large vehicles, although the confidence intervals were wide (table 2). Median air nicotine concentrations were similar in vehicles that opened windows less and more than halfway for most of the trip. However, after adjustment for the number of cigarettes, car size and other driving conditions, there was a 60% decrease in air nicotine concentrations in the vehicles that reported having their windows opened more than halfway for most of the trip (table 2).

### Knowledge and attitudes

All smokers and non-smokers acknowledged that smoking in the motor vehicle poses a health risk to the passengers (table 1). Fifty-three per cent of the smokers reported that being unable to smoke in their cars would help them to quit smoking. Ninety-three per cent of smokers agreed with the statement that motor vehicles should be smoke-free on a voluntarily basis and only 7% of the smokers agreed that motor vehicles should be smoke-free by regulation.

### DISCUSSION

In this study, conducted in motor vehicles under actual driving conditions, air nicotine concentrations measured in smokers' vehicles were strikingly higher than the non-detectable concentrations found in non-smokers' vehicles. Air nicotine concentrations within the vehicle increased with the number of cigarettes smoked. The size of the vehicle and the use of ventilation were also relevant determinants of air nicotine

concentrations, although opening windows more than halfway for most of the trip could not eliminate exposure to secondhand smoke. The location of the sampler (front seat or rear seat) had relatively little impact on nicotine concentrations within the vehicle.

Our findings are consistent with other studies showing high concentrations of particulate matter of less than  $2.5 \mu\text{m}$  in diameter ( $\text{PM}_{2.5}$ ) measured after smoking a cigarette under stationary and moving experimental conditions.<sup>7 8 14 17</sup> While ventilation rates markedly changed depending on window opening, vehicle speed and air conditioning,  $\text{PM}_{2.5}$  and carbon monoxide concentrations increased rapidly under all experimental conditions. After converting average  $\text{PM}_{2.5}$  concentration during the time over which the cigarette was smoked and its decay period to a 24-hour reference time, smoking two cigarettes within the vehicle resulted in  $\text{PM}_{2.5}$  concentrations equal to  $42 \mu\text{g}/\text{m}^3$ , higher than the 24-hour US Environmental Protection Agency's ambient air quality standard for  $\text{PM}_{2.5}$  ( $35 \mu\text{g}/\text{m}^3$ ).<sup>7</sup> With recirculation on and closed windows, peak  $\text{PM}_{2.5}$  concentrations were much higher.<sup>7</sup> In another study,  $\text{PM}_{2.5}$  concentrations after smoking a cigarette were 10-fold higher than  $\text{PM}_{2.5}$  concentrations emitted from an idling eco-diesel engine.<sup>14</sup> Studies conducted during actual driving conditions also measured high  $\text{PM}$  concentrations in Boston, US<sup>6</sup> and Crete, Greece.<sup>18</sup>

Information on smoking behaviour within cars is also available through questionnaires and observation. A cross-sectional study of smoking in cars in New Zealand found a 4.1% point prevalence of smoking in cars, with higher prevalence in lower compared to higher socioeconomic status areas, and with 23.7% of smoking cars having other occupants exposed to secondhand smoke.<sup>19</sup> In Canada, children in grades 5–12 who had been passengers in smoking cars were nearly twice as likely to respond that smoking should not be allowed in cars with children.<sup>20 21</sup> Moreover, being exposed to secondhand smoke in vehicles has been associated with an increased risk of current and persistent wheeze<sup>10</sup> and higher levels of nicotine-dependent symptoms in children,<sup>22</sup> and with increased respiratory symptoms in young adults.<sup>23</sup>



Secondhand tobacco smoke health risks have led to the introduction and adoption of legislations to protect passengers from exposure to secondhand tobacco smoke within motor vehicles. So far, legislation has focused on protecting children, with differences in the age range considered. In Arkansas, legislation restricts smoking in motor vehicles where children under age 6 are present, corresponding to the age that children are required to be in car seats.<sup>13</sup> In Louisiana and Puerto Rico, smoking bans in motor vehicles cover children up to age 13.<sup>24</sup> In California, legislation bans smoking in all vehicles with passengers under age 18.<sup>25</sup> At least 10 other US states and several countries have introduced or passed legislation to protect motor vehicle passengers from secondhand smoke exposure. Other countries such as New Zealand have invested in media campaigns advocating no smoking in motor vehicles.<sup>19</sup> Public support for smoke-free motor vehicles is also increasing, being as high as 96% of respondents in some surveys.<sup>26</sup> As more public places are becoming smoke-free, legislations in motor vehicles are less viewed as government intrusion into private matters, especially when children are included. Indeed, ethical analyses have shown that banning smoking in cars while children are present is the most ethically justifiable position, emphasising the concept of non-maleficence with regard to child passengers.<sup>27</sup>

While the main goal of banning smoking in motor vehicles is to protect non-smoking passengers from the health effects of secondhand smoke, additional benefits may include helping smokers to quit and reducing motor vehicle accidents. In our study, 53% of the smoking participants reported that banning smoking in motor vehicles could help them quit. Research conducted in Australia found that smokers had an increased risk of being involved in a motor vehicle accident than their non-smoking counterparts, and driving while smoking also increased this risk.<sup>15</sup> In Taiwan, smokers had nearly double risk of motor vehicle mortality than non-smokers after adjustment for socio-economic status and alcohol use were taken into account.<sup>28</sup> Banning smoking in cars may thus contribute to reducing car accidents. From an economic perspective, implementing smoking restrictions in cars may also prove beneficial during the sale of a vehicle. A study conducted in San Diego, California, found smoking status within a vehicle to be significantly associated with the vehicle asking price upon resale even when controlling for Kelley Blue Book (KBB) value and make of the car, reflecting changes in preferences for smoke-free environments,<sup>29</sup> as well as the residual secondhand smoke that can accumulate within the vehicles for several days after a cigarette is smoked.<sup>30</sup>

Strengths of our study include the measurement of nicotine, a specific tracer for tobacco smoke that is commonly used to assess secondhand smoke exposure as a surrogate for other toxic and carcinogenic components in tobacco.<sup>11 31</sup> Moreover, while recall bias may have affected the reporting of the number of cigarettes smoked, window opening and air conditioning use, air nicotine is an objective marker of secondhand smoke that is not affected by self-report. Our questionnaire requested information on filtered vs non-filtered cigarettes. Information on the type (that is, light vs regular) and brand of filtered cigarettes, however, was not collected and could not be related to air nicotine concentrations. Because this study aimed to measure secondhand smoke during actual driving conditions, air exchange rates of ventilation patterns within the vehicles were not directly measured. Also, uncertainties in the implementation of study procedures by the driver such as the installation and use of samplers as directed, including opening and closing the samplers, is a potential limitation of this study. While the sample size was small and multivariable models must be

### What this paper adds

- ▶ This study measured secondhand smoke exposure, as determined by air nicotine concentrations, in motor vehicles during actual driving conditions.
- ▶ Air nicotine concentrations in smokers' vehicles were strikingly higher than the non-detectable concentrations in non-smokers' vehicles and increased with the number of cigarettes smoked.
- ▶ These high levels of secondhand smoke exposure support the need for legislation that bans smoking in motor vehicles when passengers are present.

interpreted cautiously, we had sufficient statistical power to detect differences in air nicotine concentrations in smoking and non-smoking vehicles and to evaluate exposure levels with increasing number of cigarettes smoked. Finally, various types of vehicles including compact and midsize cars, minivans and SUVs were represented in the study.

While there is no safe level of secondhand smoke exposure, air nicotine concentrations in motor vehicles were generally higher than air nicotine concentrations measured in public<sup>32</sup> or private indoor places,<sup>33</sup> and even higher than concentrations measured in restaurants and bars that allow smoking.<sup>34</sup> Because smoking in cars can contaminate the vehicle surface and environment after active smoking has occurred,<sup>30</sup> further research is needed to assess possible exposure during periods following active smoking. For instance, studies comparing air nicotine concentrations during driving periods to air nicotine concentrations during 24-hour periods could be useful. Meanwhile, the high air nicotine concentrations measured in motor vehicles in this study support the urgent need for smoke-free education campaigns and legislative measures banning smoking in motor vehicles when passengers, especially children, are present. This study represents one of the few to quantify nicotine concentrations in cars and, although a pilot study, it has implications for informing larger studies and supporting policies aimed at reducing exposure to secondhand smoke in motor vehicles.

**Acknowledgements:** The authors thank Dr Jonathan Samet for his comments and suggestions provided to a previous version of this manuscript.

**Funding:** This project was supported by a Clinical Investigator Award from the Flight Attendant Medical Research Institute (FAMRI).

**Competing interests:** None.

**Provenance and peer review:** Not commissioned; externally peer reviewed.

### REFERENCES

1. **US Department of Health and Human Services.** *The health consequences of involuntary exposure to tobacco smoke: a report of the Surgeon General.* Atlanta, GA: USDHHS, 2006.
2. **Schmidt CW.** A change in the air: smoking bans gain momentum worldwide. *Environ Health Perspect* 2007;**115**:A412–5.
3. **Koh HK, Joossens LX, Connolly GN.** Making smoking history worldwide. *N Engl J Med* 2007;**356**:1496–8.
4. **Klerman L.** Protecting children: reducing their environmental tobacco smoke exposure. *Nicotine Tob Res* 2004;**6**(suppl 2):S239–53.
5. **Centers for Disease Control and Prevention.** State-specific prevalence of smoke-free home rules—United States, 1992–2003. *MMWR Morb Mort Wkly Rep* 2007;**56**:501–4.
6. **Rees VW, Connolly GN.** Measuring air quality to protect children from secondhand smoke in cars. *Am J Prev Med* 2006;**31**:363–8.
7. **Ott W, Klepeis N, Switzer P.** Air change rates of motor vehicles and in-vehicle pollutant concentrations from secondhand smoke. *J Expo Sci Environ Epidemiol* 2008;**18**:312–25.
8. **Park JH, Spengler JD, Yoon DW, et al.** Measurement of air exchange rate of stationary vehicles and estimation of in-vehicle exposure. *J Expo Anal Environ Epidemiol* 1998;**8**:65–78.

## Research paper

9. **Sendzik T**, Fong GT, Travers MJ, *et al*. An experimental investigation of tobacco smoke pollution in cars. *Nicotine Tob Res* 2009;**11**:627–34.
10. **Sly PD**, Devereill M, Kusel MM, *et al*. Exposure to environmental tobacco smoke in cars increases the risk of persistent wheeze in adolescents. *Med J Aust* 2007;**186**:322.
11. **Barnoya J**, Glantz SA. Cardiovascular effects of secondhand smoke: nearly as large as smoking. *Circulation* 2005;**111**:2684–98.
12. **Delfino RJ**, Staimer N, Tjoa T, *et al*. Personal and ambient air pollution exposures and lung function decrements in children with asthma. *Environ Health Perspect* 2008;**116**:550–8.
13. **Knapp A**. *States go after smoking in vehicles with kids*. Stateline.org; 2007. Available at: <http://stateline.org/live/printable/story?contentId=186298>. (Accessed 4 October 2008.)
14. **Invernizzi G**, Ruprecht A, Mazza R, *et al*. Particulate matter from tobacco versus diesel car exhaust: an educational perspective. *Tob Control* 2004;**13**:219–21.
15. **American Society of Heating, Refrigerating and Air-Conditioning Engineers**. *Environmental tobacco smoke: position document*. Atlanta, GA: ASHRAE, 2005.
16. **Hammond SK**, Leaderer BP. A diffusion monitor to measure exposure to passive smoking. *Environ Sci Technol* 1987;**21**:494–7.
17. **Edwards R**, Wilson N, Piersie N. Highly hazardous air quality associated with smoking in cars: New Zealand pilot study. *N Z Med J* 2006;**119**:U2294.
18. **Vardavas CI**, Linardakis M, Kafatos AG. Environmental tobacco smoke exposure in motor vehicles: a preliminary study. *Tob Control* 2006;**15**:415.
19. **Martin J**, George R, Andrews K, *et al*. Observed smoking in cars: a method and differences by socioeconomic area. *Tob Control* 2006;**15**:409–11.
20. **Leatherdale ST**, Smith P, Ahmed R. Youth exposure to smoking in the home and in cars: how often does it happen and what do youth think about it? *Tob Control* 2008;**17**:86–92.
21. **Leatherdale ST**, Ahmed R. Second-hand smoke exposure in homes and in cars among Canadian youth: current prevalence, beliefs about exposure, and changes between 2004 and 2006. *Cancer Causes Control* 2009;**20**:855–65.
22. **Belanger M**, O'Loughlin J, Okoli CT, *et al*. Nicotine dependence symptoms among young never-smokers exposed to secondhand tobacco smoke. *Addict Behav* 2008;**33**:1557–63.
23. **Lee DJ**, Dietz NA, Arheart KL, *et al*. Respiratory effects of secondhand smoke exposure among young adults residing in a "clean" indoor air state. *J Community Health* 2008;**33**:117–25.
24. **McCarthy K**. OLR research report: bans on smoking in vehicles carrying children. State of Connecticut General Assembly, ed. 2008-R-0133. 15 Feb 2008.
25. **American Lung Association**. State legislated actions on tobacco issues (SLATI) information: California; 2008. Available at: <http://slati.lungusa.org/state-templ.asp?id=5>. (Accessed 4 October 2008.)
26. **Thomson G**, Wilson N. Public attitudes to laws for smokefree private vehicles: a brief review. *Tob Control* 2008 (Epub ahead of print).
27. **Jarvie JA**, Malone RE. Children's secondhand smoke exposure in private homes and cars: an ethical analysis. *Am J Public Health* 2008;**98**:2140–5.
28. **Wen CP**, Tsai SP, Cheng TY, *et al*. Excess injury mortality among smokers: a neglected tobacco hazard. *Tob Control* 2005;**14**:i28–i32.
29. **Matt GE**, Romero R, Ma DS, *et al*. Tobacco use and asking prices of used cars: prevalence, costs, and new opportunities for changing smoking behavior. *Tob Induc Dis* 2008;**4**:2.
30. **Matt GE**, Quintana PJ, Hovell MF, *et al*. Residual tobacco smoke pollution in used cars for sale: air, dust, and surfaces. *Nicotine Tob Res* 2008;**10**:1467–75.
31. **Daisey JM**. Tracers for assessing exposure to environmental tobacco smoke: what are they tracing? *Environ Health Perspect* 1999;**107**(suppl 2):319–27.
32. **Navas-Acien A**, Peruga A, Breyse P, *et al*. Secondhand tobacco smoke in public places in Latin America, 2002–2003. *JAMA* 2004;**291**:2741–5.
33. **Wipfli H**, vila-Tang E, Navas-Acien A, *et al*. Secondhand smoke exposure among women and children: evidence from 31 countries. *Am J Public Health* 2008;**98**:672–9.
34. **Nebot M**, Lopez MJ, Gorini G, *et al*. Environmental tobacco smoke exposure in public places of European cities. *Tob Control* 2005;**14**:60–3.

## Access the latest content chosen by our Editors

BMJ Journals editors select an article from each issue to be made free online immediately on publication. Other material is free after 12 months to non-subscribers. Access the Editor's Choice from the home page—or expand your horizons and see what the other BMJ Journals editors have chosen by following the links on any BMJ Journal home page.