

Adolescent Risk Behaviors and Use of Electronic Vapor Products and Cigarettes

Zewditu Demissie, PhD,^{a,b} Sherry Everett Jones, PhD,^a Heather B. Clayton, PhD,^a Brian A. King, PhD^c

abstract

BACKGROUND: Adolescent use of tobacco in any form is unsafe; yet the use of electronic cigarettes and other electronic vapor products (EVPs) has increased in recent years among this age group. We assessed the prevalence and frequency of cigarette smoking and EVP use among high school students, and associations between health-risk behaviors and both cigarette smoking and EVP use.

METHODS: We used 2015 national Youth Risk Behavior Survey data ($N = 15\,624$) to classify students into 4 mutually exclusive categories of smoking and EVP use based on 30-day use: nonuse, cigarette smoking only, EVP use only, and dual use. Prevalence of cigarette smoking and EVP use were assessed overall and by student demographics and frequency of use. Prevalence ratios were calculated to identify associations with health risk-behaviors.

RESULTS: In 2015, 73.5% of high school students did not smoke cigarettes or use EVPs, 3.2% smoked cigarettes only, 15.8% used EVPs only, and 7.5% were dual users. Frequency of cigarette smoking and EVP use was greater among dual users than cigarette-only smokers and EVP-only users. Cigarette-only smokers, EVP-only users, and dual users were more likely than nonusers to engage in several injury, violence, and substance use behaviors; have ≥ 4 lifetime sexual partners; be currently sexually active; and drink soda ≥ 3 times/day. Only dual users were more likely than nonusers not to use a condom at last sexual intercourse.

CONCLUSIONS: EVP use, alone and concurrent with cigarette smoking, is associated with health-risk behaviors among high school students.

FREE

^aDivision of Adolescent and School Health, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention and ^bOffice on Smoking and Health, National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention, Atlanta, Georgia; and ^cUS Public Health Service Commissioned Corps, Rockville, Maryland

Dr Demissie conceptualized and designed the study, performed the data analysis, interpreted the study findings, and drafted the initial manuscript; Drs Everett Jones and Clayton conceptualized and designed the study, provided interpretation of the data, and reviewed and revised the manuscript; Dr King provided interpretation of the data and reviewed and revised the manuscript; and all authors approved the final manuscript as submitted.

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention or the US Public Health Service Commissioned Corps.

DOI: 10.1542/peds.2016-2921

Accepted for publication Nov 17, 2016

Address correspondence to Zewditu Demissie, PhD, Division of Adolescent and School Health, Centers for Disease Control and Prevention, 1600 Clifton Rd, MS E-75, Atlanta, GA 30329. E-mail: izj5@cdc.gov

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

WHAT'S KNOWN ON THIS SUBJECT: Although cigarette smoking among adolescents has decreased, the use of certain emerging tobacco products, such as electronic vapor products (EVPs), has increased. Risks, including health-risk behaviors, associated with EVP use have not yet been fully explored.

WHAT THIS STUDY ADDS: EVP use, alone and concurrent with cigarette smoking, is associated with several health-risk behaviors among high school students. This suggests comprehensive efforts to address health-risk behaviors among adolescents are warranted, including prevention strategies focused on all forms of tobacco use.

To cite: Demissie Z, Everett Jones S, Clayton HB, et al. Adolescent Risk Behaviors and Use of Electronic Vapor Products and Cigarettes. *Pediatrics*. 2017;139(2):e20162921

Cigarette smoking is the leading cause of preventable disease and death in the United States.^{1–3} Each year in the United States, an estimated 480 000 individuals die of smoking, and >3200 adolescents smoke their first cigarette each day.² Data from the National Youth Tobacco Survey indicates that cigarette smoking decreased among high school students in the United States during 2011–2015.⁴ Similarly, data from the Youth Risk Behavior Survey (YRBS) has shown that cigarette smoking among this group is at an all-time low since 1991.⁵ However, this success is tempered by the finding that use of certain emerging tobacco products, including electronic vapor products (EVPs), has increased.⁴ This has resulted in no change in overall tobacco use among middle and high school students.⁴

EVPs, also known as electronic nicotine delivery systems,⁶ include electronic cigarettes (e-cigarettes), electronic cigars, electronic pipes, vape pipes or pens, and electronic hookahs or hookah pens. EVPs are handheld devices that provide an aerosol that typically includes nicotine, additives, and other harmful and potentially harmful substances that are inhaled by the user.^{6–8} Data from the National Youth Tobacco Survey indicate that the percentage of US high school students who reported using an e-cigarette in the past 30 days increased from 1.5% to 16.0% during 2011–2015.⁴

There is a range of potential impacts, including risks and possible benefits, of EVPs on patterns of use of cigarettes and other combustible tobacco products among adults.² However, among youth, any form of tobacco use, including EVPs, is unsafe.² EVPs typically contain nicotine; nicotine exposure during adolescence can cause addiction, might harm brain development, and could lead to sustained tobacco product use.^{2,9,10} Furthermore, longitudinal studies suggest that

youth who use EVPs are more likely to subsequently initiate cigarette smoking.^{11–13} The use of multiple tobacco products among adolescents, including EVPs, is also of concern because multiple tobacco product users have greater nicotine dependence than single-product users.^{14,15} Accordingly, preventing adolescents from initiating the use of any form of tobacco product, including EVPs, is important to tobacco use prevention and control strategies.¹⁶

Given that EVPs are relatively new to the US marketplace, little is known about the use of these products in the context of other health behaviors, which can persist throughout life and contribute to significant morbidity and mortality during adolescence and adulthood.⁵ Previous research on adolescent tobacco use indicates that cigarette smoking is associated with behaviors related to injury and violence, alcohol and drug use, sexual risk behaviors, physical activity, and school problems (ie, truancy).^{17–24} Additionally, research suggests that cigarette smoking may be a marker for these and other problem behaviors.^{23,25} However, the tobacco product landscape continues to diversify, and EVPs are now the most commonly used tobacco products among US high school students, surpassing conventional cigarettes in 2014.²⁶ To date, studies on the association between EVPs and health-risk behaviors among adolescents and young adults has been limited to examining the associations between e-cigarette and substance use. These studies have found that use of e-cigarettes was associated with alcohol use, binge drinking, and marijuana use.^{12,27–31}

Several studies have assessed the prevalence and reasons for e-cigarette use; however, none have examined associated health-risk behaviors.^{13,32,33} To address this gap in the literature, this study examined the association between

EVPs and a broad range of health-risk behaviors among high school students using data from the national YRBS. This study also compares the frequency of EVP and cigarette use among exclusive and dual users. Given the considerable increases in EVP use and dual tobacco use among adolescents in recent years, the findings from this study could provide critical information to guide public health promotion efforts.

METHODS

Study Sample

The national YRBS monitors 6 priority health-risk behaviors biennially among high school students.⁵ The survey uses a 3-stage cluster sample design to obtain a nationally representative sample of public and private school students in grades 9 through 12 in the 50 states and the District of Columbia. Student participation is anonymous and voluntary, and local parental permission procedures are used. The survey is self-administered and students record their responses directly on a computer-scannable questionnaire or answer sheet. Sampling weights are applied to each record to adjust for nonresponse and the oversampling of black and Hispanic students. Additional details regarding YRBS sampling and psychometric properties have been published elsewhere.^{34,35} In 2015, the YRBS school response rate was 69%, the student response rate was 86%, the overall response rate was 60%, and the sample size was 15 624. The demographic distribution of the 2015 YRBS sample has been published elsewhere.⁵ An institutional review board at the Centers for Disease Control and Prevention approved the national YRBS.

Measures

Cigarette Smoking and EVP Use

Current cigarette smoking and EVP use were assessed with the following

2 questions: "During the past 30 days, on how many days did you smoke cigarettes?" and "During the past 30 days, on how many days did you use an electronic vapor product?" An introduction to the EVP question was provided to give examples of EVP brands (blu, NJOY, or Starbuzz) and examples of EVP types (e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens). Response options for both questions were as follows: 0 days, 1 to 2 days, 3 to 5 days, 6 to 9 days, 10 to 19 days, 20 to 29 days, and all 30 days. These questions were examined in 2 ways. First, each question was used in its original form to examine the number of days students smoked cigarettes and used EVPs (ie, frequency of use). Second, they were combined to create a 4-level variable with mutually exclusive categories: nonuse, cigarette smoking only, EVP use only, and dual use (cigarettes and EVP).

Health-Risk Behaviors

Dependent variables included those related to unintentional injuries and violence, substance use, and sexual risk behaviors (Table 1). Variables in these behavioral domains were included in a previous YRBS analysis that examined associations between tobacco use and health-risk behaviors.²¹ The current study also assessed dietary behaviors and physical activity (Table 1). The cut points used for fruit and vegetable consumption were based on earlier fruit and vegetable recommendations that used times/day as the measurement unit³⁶ because current fruit and vegetable intake recommendations are reported in cup equivalents/day,³⁷ which cannot be calculated using YRBS data. The soda cutpoint was based on a study of Americans aged 2 years and older, which found that the estimated 90th percentile of daily energy intake from sugar-sweetened beverages was 450 kcal (equivalent to three 12-oz cans of soda).³⁸ The physical activity

cutpoint was based on national recommendations.³⁹

Covariates

Three demographic characteristics were included in this analysis: sex (male, female), grade (9th, 10th, 11th, and 12th), and race/ethnicity. Students were classified into 4 racial/ethnic categories: white, non-Hispanic; black, non-Hispanic; Hispanic or Latino (of any race); and other or multiple race. The numbers of students in the other or multiple racial/ethnic group were too small for meaningful analysis. The YRBS questionnaire includes questions about 2 other types of tobacco use. Students were also asked their current (past 30-day) use of (1) smokeless tobacco products (chewing tobacco, snuff, or dip) as well as (2) cigars, cigarillos, or little cigars. The response options for these questions were split into 2 groups: any use and no use.

Statistical Analysis

Analyses were conducted on weighted data using SUDAAN (Research Triangle Park, NC), a software package that accounts for the survey's complex sampling design. The percentage of students in each category of the 4-level combined cigarette and EVP variable was calculated among students overall and by sex, race/ethnicity, and grade. χ^2 tests were conducted to assess differences by demographics. Among cigarette users, the frequency of cigarette use was compared between cigarette-only smokers and dual users using pairwise comparisons (*t* statistic). In parallel, among EVP users, the frequency of EVP use was compared between EVP only and dual users using pairwise comparisons. Additionally, multivariable logistic regression models adjusted for sex, race/ethnicity, grade, smokeless tobacco use, and cigar smoking were used to estimate adjusted prevalence ratios (PRs) and 95% confidence intervals (CIs) for the associations

between cigarette smoking or EVP use and health-risk behaviors using nonusers as the referent group. PRs were considered statistically significant if 95% CIs did not include 1.0. Lastly, linear contrasts were conducted to compare results for all health-risk behaviors by cigarette and EVP group using other referents (EVP only vs cigarette only, dual use vs cigarette only, and dual use vs EVP only). Differences were considered significant if the *P* value was < .05.

RESULTS

Prevalence of Use

Nationwide, 73.5% of high school students did not smoke cigarettes or use EVPs, 3.2% smoked cigarettes only, 15.8% used EVPs only, and 7.5% were dual users of both cigarettes and EVPs (Table 2). The distribution across the 4-level cigarette and EVP use variable differed significantly by sex (*P* = .04), race/ethnicity (*P* < .001), and grade level (*P* < .001).

Frequency of Use

Dual users of cigarettes and EVPs were significantly less likely than cigarette-only smokers to smoke cigarettes 1 to 2 days (32.8% vs 46.0%) but significantly more likely to smoke cigarettes 10 to 19 days (10.7% vs 5.3%) and 20 to 29 days (12.3% vs 4.1%) (Fig 1). Dual users of cigarettes and EVPs were significantly less likely than EVP-only users to use EVPs 1 to 2 days (29.2% vs 55.6%), but significantly more likely to use EVPs 6 to 9 days (14.7% vs 9.1%), 10 to 19 days (15.3% vs 7.5%), 20 to 29 days (5.8% vs 3.1%), and all 30 days (13.0% vs 5.3%) (Fig 2).

Cigarettes, EVP, and Health-Risk Behaviors

Unintentional Injuries and Violence

Cigarette-only smokers, EVP-only users, and dual users were significantly more likely than

TABLE 1 Question Wording and Analytic Coding for Included Health Risk Behaviors, National YRBS, 2015

Health Risk Behavior	Questionnaire Item	Analytic Coding
Injury and violence		
Text/e-mail while driving	During the past 30 days, on how many days did you text or e-mail while driving a car or other vehicle? ^a	≥1 vs 0 days
Engaged in a physical fight	During the past 12 months, how many times were you in a physical fight?	≥1 vs 0 times
Carried a weapon	During the past 30 days, on how many days did you carry a weapon such as a gun, knife, or club?	≥1 vs 0 days
Attempted suicide	During the past 12 months, how many times did you actually attempt suicide?	≥1 vs 0 times
Substance use		
Current alcohol use	During the past 30 days, on how many days did you have at least 1 drink of alcohol?	≥1 vs 0 days
Current marijuana use	During the past 30 days, how many times did you use marijuana?	≥1 vs 0 times
Ever synthetic marijuana use	During your life, how many times have you used synthetic marijuana (also called K2, Spice, fake weed, King Kong, Yucatan Fire, Skunk, or Moon Rocks)?	≥1 vs 0 times
Ever use of other illicit drugs	Combines all of the following individual questionnaire items: During your life, how many times have you used any form of cocaine, including powder, crack, or freebase? During your life, how many times have you snuffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high? During your life, how many times have you used heroin (also called smack, junk, or China White)? During your life, how many times have you used methamphetamines (also called speed, crystal, crank or ice)? During your life, how many times have you used ecstasy (also called MDMA)? During your life, how many times have you used hallucinogenic drugs, such as LSD, acid, PCP, angel dust, mescaline, or mushrooms?	≥1 vs 0 times
Ever nonmedical use of prescription drugs	During your life, how many times have you taken a prescription drug (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax) without a doctor's prescription?	≥1 vs 0 times
Sexual risk behaviors		
Lifetime sexual partners	During your life, with how many people have you had sexual intercourse?	≥4 vs <4 persons
Currently sexually active	During the past 3 months, with how many people did you have sexual intercourse?	≥1 vs 0 persons
Condom use at last sexual intercourse	The last time you had sexual intercourse, did you or your partner use a condom? ^b	No vs yes
Dietary behaviors and physical activity		
Fruit intake	Combines 2 questionnaire items: During the past 7 days, how many times did you eat fruit? (Do not count fruit juice.) During the past 7 days, how many times did you drink 100% fruit juices such as orange juice, apple juice, or grape juice? (Do not count punch, Kool-Aid, sports drinks, or other fruit-flavored drinks.)	<2 times/day vs ≥2 times/day
Vegetable intake	Combines 4 questionnaire items: During the past 7 days, how many times did you eat green salad? During the past 7 days, how many times did you eat potatoes? (Do not count French fries, fried potatoes, or potato chips.) During the past 7 days, how many times did you eat carrots? During the past 7 days, how many times did you eat other vegetables? (Do not count green salad, potatoes, or carrots.)	<3 times/day vs ≥3 times/day
Soda intake	During the past 7 days, how many times did you drink a can, bottle, or glass of a sports drink such as Gatorade or PowerAde? (Do not count low-calorie sports drinks such as Propel or G2.)	≥3 times/day vs <3 times/day
Physical activity	During the past 7 days, on how many days were you physically active for a total of at least 60 minutes per day? (Add up all the time you spent in any kind of physical activity that increased your heart rate and made you breathe hard some of the time.)	<7 days vs 7 days

^a Excludes students who did not drive a car or other vehicle during the past 30 days.^b Excludes students who were not currently sexually active.

nonusers to engage in a physical fight (PR range: 1.72–2.87) and attempt suicide (PR range: 1.86–4.01) (Table 3). Both EVP-only users and dual users were significantly more likely than nonusers to text or e-mail while driving (PR: 1.39 and 1.41, respectively). Only dual users were significantly more likely than nonusers to carry a weapon (PR: 1.82; 95% CI: 1.30–2.56).

Substance Use Behaviors

Cigarette-only smokers, EVP-only users, and dual users were significantly more likely than nonusers to currently drink alcohol (PR range: 2.62–3.29), currently use marijuana (PR range: 3.49–5.22), ever use synthetic marijuana (PR range: 3.67–8.65), ever use other illicit drugs (PR range: 2.73–5.75),


and report ever nonmedical use of prescription drugs (PR range: 2.30–4.13) (Table 3).

Sexual Risk Behaviors

Cigarette-only smokers, EVP-only users, and dual users were significantly more likely than nonusers to have ≥4 lifetime partners (PR range: 2.35–4.60) and be currently sexually active (PR

TABLE 2 Prevalence (95% CI) of Cigarette Use and Electronic Vapor Product Use, National YRBS, 2015

Demographic	Nonuse	Cigarette Only	EVP Only	Dual Use	$\chi^2 (P)$
Overall	73.5 (71.1–75.8)	3.2 (2.6–3.9)	15.8 (14.2–17.5)	7.5 (6.5–8.8)	
Sex					3.1 (.04)
Female	75.0 (72.3–77.5)	2.9 (2.2–3.9)	15.4 (13.8–17.1)	6.7 (5.6–8.1)	
Male	72.0 (69.0–74.8)	3.5 (2.8–4.4)	16.3 (14.1–18.7)	8.3 (7.1–9.6)	
Race/ethnicity					5.5 (<.001)
White, non-Hispanic	71.9 (68.2–75.4)	3.6 (2.7–4.6)	15.6 (13.7–17.7)	8.9 (7.1–11.1)	
Black, non-Hispanic	80.4 (76.1–84.1)	2.9 (1.7–4.9)	13.6 (11.0–16.7)	3.1 (2.1–4.5)	
Hispanic	71.6 (68.9–74.2)	2.8 (2.1–3.6)	19.2 (17.0–21.6)	6.4 (5.4–7.6)	
Grade					14.8 (<.001)
9	79.1 (76.5–81.5)	1.8 (1.2–2.7)	13.4 (11.2–15.9)	5.7 (4.2–7.7)	
10	75.2 (71.6–78.4)	2.3 (1.7–3.1)	16.1 (13.9–18.5)	6.4 (5.0–8.2)	
11	70.5 (67.2–73.5)	4.4 (3.4–5.7)	16.6 (14.1–19.5)	8.5 (6.9–10.3)	
12	68.4 (64.6–71.9)	4.4 (3.3–6.0)	17.6 (15.1–20.4)	9.6 (7.8–11.7)	

**FIGURE 1**

Cigarette smoking frequency among cigarette only smokers and dual users, National YRBS, 2015. "Dual user" is defined as a student who currently smokes cigarettes and uses electronic vapor products. *Statistical difference between cigarette-only smokers and dual users.

range: 1.86–2.30) (Table 3). Only dual users were significantly more likely than nonusers not to use a condom at last sexual intercourse (PR: 1.27; 95% CI: 1.07–1.50).

Dietary Behaviors and Physical Activity

Dual users were significantly less likely to eat vegetables <3 times per day compared with nonusers (PR: 0.94; 95% CI: 0.91–0.98) (Table 3). Cigarette-only smokers, EVP-only users, and dual users were significantly more likely than nonusers to drink soda ≥3 times/day (PR range: 1.35–3.03). EVP-only users were significantly less likely not to engage in daily physical

activity compared with nonusers (PR: 0.91; 95% CI: 0.88–0.95).

Linear Contrasts

Engaging in health-risk behaviors did not generally differ between EVP-only users and cigarette-only smokers (Table 3); however, cigarette-only smokers were significantly more likely than EVP-only users to attempt suicide, ever use synthetic marijuana, have ≥4 lifetime sexual partners, drink soda ≥3 times/day, and be physically active <7 days in the 7 days before the survey. When comparing dual users to cigarette-only smokers, dual users were significantly more likely than cigarette-only smokers

to engage in all substance use behaviors and physical fighting. When comparing dual users to EVP-only users, significant differences were observed for all substance use and sexual risk behaviors, physical fighting, attempted suicide, soda intake, and physical activity. In all cases, dual users were more likely to engage in these behaviors compared with EVP-only users.

DISCUSSION

The findings of this study are consistent with other studies documenting an association between EVP use, particularly e-cigarettes, and other substance use.^{12,27–31} However, this study also shows that EVP use is associated with other health-risk behaviors. For a majority of the behaviors examined in this study, cigarette-only smokers and EVP-only users were as likely to engage in that behavior, suggesting that the associations between health-risk behaviors and EVP use mirror the associations between health-risk behaviors and conventional cigarettes. The pattern of cigarette-only and EVP-only users having intermediate risks between nonusers and dual users is consistent with other published studies.^{29,30} Consequently, addressing the diversity of tobacco use as part of youth health promotion efforts is important.




FIGURE 2

EVP use frequency among EVP-only users and dual users, National YRBS, 2015. “Dual user” is defined as a student who currently smokes cigarettes and uses electronic vapor products. *Statistical difference between EVP only users and dual users.

Research on conventional cigarettes suggests that smoking occurs along with, or clusters with, various health-risk behaviors and that cigarette smoking may serve as a marker for these behaviors.^{17–25} This is important to consider in the context of public health practice because research indicates that among adults, the odds of reporting fair or poor health status increase with each unhealthy behavior that an individual engages in.⁴⁰ The development of certain unhealthy behaviors during adolescence puts adolescents on trajectories toward chronic disease.²⁵ The reason for the association between tobacco use and other health-risk behaviors may be due to common factors (ie, lack of resources such as health education), motivation by similar goals (ie, rejecting conventional social norms), or the activation of similar neural pathways.^{24,25,41} Cigarette smoking has also been found to be associated with emotional and behavioral problems, which could also make adolescents more vulnerable to developing other health-risk behaviors.^{42,43} Future studies may show similar vulnerabilities among EVP users, including whether e-cigarette use may be introducing

youth with relatively low-risk profiles to substance use. Therefore, comprehensive efforts to address the health-risk behaviors among adolescents are warranted, including prevention strategies focused on all forms of tobacco use, including EVPs.

Despite the harms of EVP use among adolescents, use of these products among US middle and high school students has increased considerably in recent years, for several possible reasons. First, in 2014, 18 million youth were exposed to e-cigarette advertising.⁴⁴ Given the link between marketing of conventional tobacco products and tobacco initiation among youth, it is likely that increases in e-cigarette marketing are contributing to increases in EVP use.⁴⁴ Second, the flavorings frequently used in EVPs are appealing to adolescents.^{6,8,45} Third, use of EVPs may be considered more socially acceptable, particularly in indoor areas where conventional cigarette smoking is prohibited.⁶ Fourth, EVPs have been relatively easy for youth to access; they are available in retail stores, mall kiosks, and the Internet.^{6,46} However, efforts are underway to reduce potential

public harms of these products through regulation. In May 2016, the US Food and Drug Administration finalized a rule extending its authority to all products that meet the definition of a tobacco product, including e-cigarettes. E-cigarettes are considered a tobacco product due to a 2011 court ruling.⁴⁷ The 2016 rule sets a national minimum age for sales, requires tobacco product ingredient reporting and reporting of harmful and potentially harmful constituents, and ensures Food and Drug Administration premarket review of new and changed tobacco products.⁴⁸

This study found that dual users smoked cigarettes more frequently than cigarette-only smokers and used EVPs more frequently than EVP-only users. These findings are consistent with research showing that among US adolescents who ever smoked cigarettes, e-cigarette users were less likely than nonusers to abstain from cigarettes.⁴⁹ Additionally, a study of Polish adolescents found that dual users of electronic and conventional cigarettes were more likely to smoke cigarettes daily and less likely to smoke fewer cigarettes per day than students who exclusively smoked conventional cigarettes.⁵⁰ This lack of decline in smoking frequency among youth e-cigarette users is contrary to studies showing that EVP use may reduce cigarette smoking frequency among adults.⁵¹ These findings underscore the importance of evidence-based measures to prevent use of all forms of tobacco use, including experimental or intermittent use, among youth.⁵² Studies have shown that symptoms of tobacco dependence are evident even among adolescents who used only a single tobacco product on as few as 1 to 2 days during the previous month.¹⁵

In addition to population-based interventions, such as youth access restrictions and comprehensive smoke-free policies that include

TABLE 3 Risk of Health Risk Behaviors by Cigarette Smoking and EVP Use, National YRBS, 2015

Health Risk Behavior	%	Nonuse	Cigarette Smoking Only			EVP Use Only		Dual Use ^a
			PR	%	PR (95% CI)	%	PR (95% CI)	
Injury and violence								
Text/e-mail while driving ^b	34.4	Ref	53.4	1.19 (0.94–1.49)	52.7	1.39 (1.26–1.53)	62.7	1.41 (1.21–1.64)
Engaged in a physical fight ^c	16.5	Ref	36.3	1.88 (1.52–2.33)	29.6	1.72 (1.53–1.93)	55.2 ^{d,e}	2.87 (2.53–3.26)
Carried a weapon ^b	2.5	Ref	8.0	1.48 (0.84–2.59)	4.9	1.45 (0.99–2.12)	12.0	1.82 (1.30–2.56)
Attempted suicide ^c	5.7	Ref	18.5	3.46 (2.51–4.77)	10.5 ^f	1.86 (1.53–2.25)	23.3 ^e	4.01 (3.17–5.08)
Substance use								
Current alcohol use ^b	18.7	Ref	72.7	2.67 (2.22–3.20)	61.9	2.62 (2.39–2.87)	86.4 ^{d,e}	3.29 (2.88–3.77)
Current marijuana use ^b	9.9	Ref	51.4	3.49 (2.80–4.36)	43.7	3.70 (3.16–4.32)	69.7 ^{d,e}	5.22 (4.42–6.18)
Ever synthetic marijuana use	3.2	Ref	26.1	5.39 (3.84–7.58)	14.5 ^f	3.67 (3.04–4.44)	40.2 ^{d,e}	8.65 (6.60–11.35)
Ever use of other illicit drugs ^g	6.1	Ref	29.2	3.23 (2.17–4.83)	19.8	2.73 (2.25–3.30)	50.7 ^{d,e}	5.75 (4.48–7.40)
Ever nonmedical use of prescription drugs	9.3	Ref	36.1	2.66 (2.04–3.48)	24.8	2.30 (1.97–2.68)	54.9 ^{d,e}	4.13 (3.44–4.96)
Sexual risk behaviors								
Four or more lifetime sexual partners	6.1	Ref	33.8	3.84 (2.86–5.16)	16.9 ^f	2.35 (1.93–2.87)	39.0 ^e	4.60 (3.64–5.81)
Current sexual activity ^h	21.5	Ref	56.3	1.93 (1.64–2.28)	46.2	1.86 (1.65–2.10)	63.8 ^e	2.30 (1.91–2.78)
Did not use a condom at last sexual intercourse ⁱ	41.5	Ref	48.4	1.17 (0.93–1.49)	38.1	0.93 (0.82–1.05)	52.2 ^e	1.27 (1.07–1.50)
Dietary behaviors and physical activity^j								
Fruit <2 times/d	69.4	Ref	71.5	1.05 (0.96–1.14)	65.7	0.95 (0.90–1.01)	66.8	0.99 (0.93–1.06)
Vegetables <3 times/d	86.6	Ref	82.8	0.98 (0.94–1.02)	84.5	0.98 (0.95–1.01)	79.0	0.94 (0.91–0.98)
Soda ≥3 times/d	5.3	Ref	15.3	2.66 (1.79–3.95)	7.6 ^f	1.35 (1.04–1.76)	18.1 ^e	3.03 (2.26–4.05)
<7 d of physical activity	74.1	Ref	76.4	1.07 (1.00–1.15)	66.7 ^f	0.91 (0.88–0.95)	69.5 ^e	1.01 (0.96–1.07)

PRs (adjusted for sex, race/ethnicity, grade, smokeless tobacco use, and cigar smoking): comparisons are considered significant if 95% CIs do not include 1.0. Linear contrasts: differences are considered significant if $P < .05$ for the contrast. Significant differences are noted on the prevalence estimates.

^a "Dual use" is defined as currently smoking cigarettes and using EVPs.

^b During the 30 d before the survey.

^c During the 12 mo before the survey.

^d Linear contrast is significant for dual use vs cigarette smoking only.

^e Linear contrast is significant for dual use vs EVP use only.

^f Linear contrast is significant for EVP use only vs cigarette smoking only.

^g Cocaine, inhalants, heroin, methamphetamines, ecstasy, and hallucinogenic drugs.

^h During the 3 mo before the survey.

ⁱ Analyzed among students who were currently sexually active.

^j During the 7 d before the survey.

EVPs,⁵³ actions can also be taken in the clinical setting to address youth tobacco use. Screening of patients and their caregivers for EVP use can be incorporated into tobacco use screening.⁶ Counseling about the harms associated with adolescent tobacco use, including EVP use, is critical.⁶ It is particularly important that this counseling identify and correct misperceptions about the use of EVP products in this population.⁴⁰

This study has at least 4 limitations. First, YRBS data are self-reported, and students may misreport their

behaviors. Although the EVP question is new to YRBS, a psychometric study has shown that the YRBS questions generally have good test-retest reliability.³⁴ Second, the data are from an observational study; therefore, causality cannot be determined. Third, some of the behaviors investigated may not be risky for particular individuals under certain circumstances. For example, sexual intercourse without a condom may not be risky for mutually monogamous partners when another contraceptive method is used. Lastly, data are only collected

on adolescents who attend school and therefore are not representative of all individuals in this age group. However, nationwide in 2012, ~3% of persons aged 16 to 17 years were not enrolled in a high school program and had not completed high school.⁵⁴

CONCLUSIONS

EVP use, alone and concurrent with cigarette smoking, is associated with health-risk behaviors among high school students. Moreover, dual users of EVPs and conventional

cigarettes reported using these products more frequently than exclusive users of these products. These findings underscore the importance of comprehensive efforts to address health-risk behaviors among adolescents, including

prevention strategies focused on all forms of tobacco use, including EVPs. Additionally, educational and counseling efforts focusing on the harms associated with adolescent tobacco use, including EVPs, are critical.

ABBREVIATIONS

- CI: confidence interval
EVP: electronic vapor product
PR: prevalence ratio
YRBS: Youth Risk Behavior Survey

Copyright © 2017 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

REFERENCES

- Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Actual causes of death in the United States, 2000. *JAMA*. 2004;291(10):1238–1245
- US Department of Health and Human Services. *The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General*. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2014
- US Department of Health and Human Services. *How Tobacco Smoke Causes Disease: What It Means to You*. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2010
- Singh T, Arrazola RA, Corey CG, et al. Tobacco use among middle and high school students—United States, 2011–2015. *MMWR Morb Mortal Wkly Rep*. 2016;65(14):361–367
- Kann L, McManus T, Harris WA, et al; Centers for Disease Control and Prevention. Youth risk behavior surveillance—United States, 2015. *MMWR Surveill Summ*. 2016;65(6 No. SS-6):1–174
- Walley SC, Jenssen BP; Section on Tobacco Control. Electronic nicotine delivery systems. *Pediatrics*. 2015;136(5):1018–1026
- McCarthy M. E-cigarettes could addict a new generation of youth to nicotine, doctors are told. *BMJ*. 2015;351:h5728
- Lauterstein D, Hoshino R, Gordon T, Watkins BX, Weitzman M, Zelikoff J. The changing face of tobacco use among United States youth. *Curr Drug Abuse Rev*. 2014;7(1):29–43
- US Department of Health and Human Services. *The Health Consequences of Smoking: Nicotine Addiction. A Report of the Surgeon General*. Rockville, MD: US Department of Health and Human Services, Public Health Service, Centers for Disease Control, Center for Health Promotion and Education, Office on Smoking and Health; 1988
- England LJ, Bunnell RE, Pechacek TF, Tong VT, McAfee TA. Nicotine and the developing human: a neglected element in the electronic cigarette debate. *Am J Prev Med*. 2015;49(2):286–293
- Leventhal AM, Strong DR, Kirkpatrick MG, et al. Association of electronic cigarette use with initiation of combustible tobacco product smoking in early adolescence. *JAMA*. 2015;314(7):700–707
- Unger JB, Soto DW, Leventhal A. E-cigarette use and subsequent cigarette and marijuana use among Hispanic young adults. *Drug Alcohol Depend*. 2016;163:261–264
- Barrington-Trimis JL, Urman R, Berhane K, et al. E-cigarettes and future cigarette use. *Pediatrics*. 2016;138(1):e20160379
- Lanza ST, Vasilenko SA. New methods shed light on age of onset as a risk factor for nicotine dependence. *Addict Behav*. 2015;50:161–164
- Apelberg BJ, Corey CG, Hoffman AC, et al. Symptoms of tobacco dependence among middle and high school tobacco users: results from the 2012 National Youth Tobacco Survey. *Am J Prev Med*. 2014;47(2 Suppl 1):S4–S14
- CDC Office on Smoking and Health. E-cigarette information. Available at: www.cdc.gov/tobacco/stateandcommunity/pdfs/cdc-osh-information-on-e-cigarettes-november-2015.pdf. Accessed April 14, 2016
- Cavazos-Rehg PA, Krauss MJ, Spitznagel EL, Grucza RA, Bierut LJ. Youth tobacco use type and associations with substance use disorders. *Addiction*. 2014;109(8):1371–1380
- Dunn MS. Association between physical activity and substance use behaviors among high school students participating in the 2009 Youth Risk Behavior Survey. *Psychol Rep*. 2014;114(3):675–685
- Gart R, Kelly S. How illegal drug use, alcohol use, tobacco use, and depressive symptoms affect adolescent suicidal ideation: A secondary analysis of the 2011 Youth Risk Behavior Survey. *Issues Ment Health Nurs*. 2015;36(8):614–620
- Everett SA, Giovino GA, Warren CW, Crossett L, Kann L. Other substance use among high school students

- who use tobacco. *J Adolesc Health*. 1998;23(5):289–296
21. Everett SA, Malarcher AM, Sharp DJ, Husten CG, Giovino GA. Relationship between cigarette, smokeless tobacco, and cigar use, and other health risk behaviors among U.S. high school students. *J Sch Health*. 2000;70(6):234–240
 22. Escobedo LG, Reddy M, DuRant RH. Relationship between cigarette smoking and health risk and problem behaviors among US adolescents. *Arch Pediatr Adolesc Med*. 1997;151(1):66–71
 23. Austin G, McCarthy W, Slade S, Bailet W. *Links Between Smoking and Substance Use, Violence, and Social Problems (CHKS Factsheet #5)*. Los Alamitos, CA: WestEd; 2007
 24. Pesa JA. The association between smoking and unhealthy behaviors among a national sample of Mexican-American adolescents. *J Sch Health*. 1998;68(9):376–380
 25. Spring B, Moller AC, Coons MJ. Multiple health behaviours: overview and implications. *J Public Health (Oxf)*. 2012;34(suppl 1):i3–i10
 26. Arrazola RA, Singh T, Corey CG, et al; Centers for Disease Control and Prevention (CDC). Tobacco use among middle and high school students - United States, 2011-2014. *MMWR Morb Mortal Wkly Rep*. 2015;64(14):381–385
 27. Camenga DR, Kong G, Cavallo DA, et al. Alternate tobacco product and drug use among adolescents who use electronic cigarettes, cigarettes only, and never smokers. *J Adolesc Health*. 2014;55(4):588–591
 28. Kristjansson AL, Mann MJ, Sigfusdottir ID. Licit and illicit substance use by adolescent e-cigarette users compared with conventional cigarette smokers, dual users, and nonusers. *J Adolesc Health*. 2015;57(5):562–564
 29. Wills TA, Knight R, Williams RJ, Pagano I, Sargent JD. Risk factors for exclusive e-cigarette use and dual e-cigarette use and tobacco use in adolescents. *Pediatrics*. 2015;135(1). Available at: www.pediatrics.org/cgi/content/full/135/1/e43
 30. Surís JC, Berchtold A, Akre C. Reasons to use e-cigarettes and associations with other substances among adolescents in Switzerland. *Drug Alcohol Depend*. 2015;153:140–144
 31. Saddleson ML, Kozlowski LT, Giovino GA, et al. Risky behaviors, e-cigarette use and susceptibility of use among college students. *Drug Alcohol Depend*. 2015;149:25–30
 32. Bold KW, Kong G, Cavallo DA, Camenga DR, Krishnan-Sarin S. Reasons for trying e-cigarettes and risk of continued use. *Pediatrics*. 2016;138(3):e20160895
 33. Kamboj A, Spiller HA, Casavant MJ, Chounthirath T, Smith GA. Pediatric exposure to e-cigarettes, nicotine, and tobacco products in the United States. *Pediatrics*. 2016;137(6):e20160041
 34. Brener ND, Kann L, McManus T, Kinchen SA, Sundberg EC, Ross JG. Reliability of the 1999 youth risk behavior survey questionnaire. *J Adolesc Health*. 2002;31(4):336–342
 35. Brener ND, Kann L, Shanklin S, et al; Centers for Disease Control and Prevention (CDC); Centers for Disease Control and Prevention (CDC). Methodology of the Youth Risk Behavior Surveillance System—2013. *MMWR Recomm Rep*. 2013;62(RR-1):1–20
 36. US Department of Health and Human Services. *Healthy People 2010: Understanding and Improving Health*. 2nd ed. Washington, DC: US Government Printing Office; 2000
 37. US Department of Health and Human Services and US Department of Agriculture. *2015–2020 Dietary Guidelines for Americans*. 8th ed. December 2015. Available at: <https://health.gov/dietaryguidelines/2015/guidelines>. Accessed December 3, 2016
 38. Ogden CL, Kit BK, Carroll MD, Park S. Consumption of sugar drinks in the United States, 2005–2008. *NCHS Data Brief*. 2011;71(71):1–8
 39. US Department of Health and Human Services. *2008 Physical Activity Guidelines for Americans*. Washington, DC: US Department of Health and Human Services; 2008
 40. United Health Foundation. *America's Health Rankings Spotlight: Impact of Unhealthy Behaviors*. Minnetonka, MN: United Health Foundation; 2016
 41. Turbin MS, Jessor R, Costa FM. Adolescent cigarette smoking: health-related behavior or normative transgression? *Prev Sci*. 2000;1(3):115–124
 42. Vega WA, Chen KW, Williams J. Smoking, drugs, and other behavioral health problems among multiethnic adolescents in the NHSDA. *Addict Behav*. 2007;32(9):1949–1956
 43. Giannakopoulos G, Tzavara C, Dimitrakaki C, Kolaitis G, Rotsika V, Tountas Y. Emotional, behavioural problems and cigarette smoking in adolescence: findings of a Greek cross-sectional study. *BMC Public Health*. 2010;10:57
 44. Centers for Disease Control and Prevention. E-cigarette ads and youth. Available at: www.cdc.gov/vitalsigns/ecigarette-ads/index.html. Accessed April 15, 2016
 45. Hildick-Smith GJ, Pesko MF, Shearer L, et al. A practitioner's guide to electronic cigarettes in the adolescent population. *J Adolesc Health*. 2015;57(6):574–579
 46. Rose SW, Barker DC, D'Angelo H, et al. The availability of electronic cigarettes in U.S. retail outlets, 2012: results of two national studies. *Tob Control*. 2014;23(suppl 3):iii10–iii16
 47. Sotera, Inc. v. Food & Drug Administration. Available at: www.wlf.org/Upload/litigation/briefs/SmokingEverywherevFDA-WLFAmicus.pdf. Accessed July 21, 2016
 48. Food and Drug Administration. Deeming Tobacco Products to Be Subject to the Federal Food, Drug, and Cosmetic Act, as Amended by the Family Smoking Prevention and Tobacco Control Act; Regulations on the Sale and Distribution of Tobacco Products and Required Warning Statements for Tobacco Products; Final Rule. *Fed Regist*. 2016;81(90):28973–29106
 49. Dutra LM, Glantz SA. Electronic cigarettes and conventional cigarette use among U.S. adolescents: a cross-sectional study. *JAMA Pediatr*. 2014;168(7):610–617
 50. Goniewicz ML, Leigh NJ, Gawron M, et al. Dual use of electronic and tobacco cigarettes among adolescents: a

- cross-sectional study in Poland. *Int J Public Health*. 2016;61(2):189–197
51. Adkison SE, O'Connor RJ, Bansal-Travers M, et al. Electronic nicotine delivery systems: international tobacco control four-country survey. *Am J Prev Med*. 2013;44(3):207–215
52. US Department of Health and Human Services. *Preventing Tobacco Use Among Youth and Young Adults: A Report of the Surgeon General*. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention; 2012
53. CDC Office on Smoking and Health. Electronic nicotine delivery systems: Key facts. 2015. Available at: www.cdc.gov/tobacco/stateandcommunity/pdfs/ends-key-facts2015.pdf. Accessed April 14, 2016
54. Stark P, Noel AM. *Trends in High School Dropout and Completion Rates in the United States: 1972–2012 (Publication No. NCES 2015-015)*. Washington, DC: US Department of Education, National Center for Education Statistics; 2015

Adolescent Risk Behaviors and Use of Electronic Vapor Products and Cigarettes

Zewditu Demissie, Sherry Everett Jones, Heather B. Clayton and Brian A. King

Pediatrics 2017;139;; originally published online January 23, 2017;

DOI: 10.1542/peds.2016-2921

Updated Information & Services	including high resolution figures, can be found at: /content/139/2/e20162921.full.html
References	This article cites 40 articles, 9 of which can be accessed free at: /content/139/2/e20162921.full.html#ref-list-1
Citations	This article has been cited by 3 HighWire-hosted articles: /content/139/2/e20162921.full.html#related-urls
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Adolescent Health/Medicine /cgi/collection/adolescent_health:medicine_sub Substance Abuse /cgi/collection/substance_abuse_sub Smoking /cgi/collection/smoking_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: /site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: /site/misc/reprints.xhtml

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS[®]

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Adolescent Risk Behaviors and Use of Electronic Vapor Products and Cigarettes

Zewditu Demissie, Sherry Everett Jones, Heather B. Clayton and Brian A. King

Pediatrics 2017;139;; originally published online January 23, 2017;

DOI: 10.1542/peds.2016-2921

The online version of this article, along with updated information and services, is located on the World Wide Web at:

[/content/139/2/e20162921.full.html](https://doi.org/10.1542/peds.2016-2921)

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2017 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

