

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.

## **E-cigarettes and their impact on health: from pharmacology to clinical implications**

**Authors:** Andrzej Sobczak, Leon Kosmider, Bartosz Koszowski, Maciej Ł. Goniewicz

**Article type:** Review article

**Received:** March 3, 2020.

**Accepted:** March 6, 2020.

**Published online:** March 10, 2020.

**ISSN:** 1897-9483

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License ([CC BY-NC-SA4.0](https://creativecommons.org/licenses/by-nc-sa/4.0/)), allowing third parties to copy and redistribute the material in any medium or format and to remix, transform, and build upon the material, provided the original work is properly cited, distributed under the same license, and used for noncommercial purposes only. For commercial use, please contact the journal office at [pamw@mp.pl](mailto:pamw@mp.pl).

Title: E-cigarettes and their impact on health: from pharmacology to clinical implications

Andrzej Sobczak<sup>1</sup>, Leon Kosmider<sup>1</sup>, Bartosz Koszowski<sup>2</sup>, Maciej Ł.Goniewicz<sup>3</sup>

<sup>1</sup> Department of General and Analytical Chemistry, Faculty of Pharmaceutical Sciences in Sosnowiec, Medical University of Silesia, Katowice, Poland

<sup>2</sup> Battelle Memorial Institute, Baltimore, MD 21212, USA

<sup>3</sup> Department of Health Behavior, Roswell Park Comprehensive Cancer Center, Buffalo, NY 14263 USA

Short title: E-cigarettes and clinical issues

Corresponding author: Prof. Andrzej Sobczak, PhD, Department of General and Analytical Chemistry, Faculty of Pharmaceutical Sciences in Sosnowiec, Medical University of Silesia, Jagiellońska 4, 41-200 Sosnowiec, Poland, tel. +48 32 3641564, [asobczak@sum.edu.pl](mailto:asobczak@sum.edu.pl)

Conflict of interest:

In 2015 AS accepted personal fees from the eSmoking Institute in Poznań, Poland, and non-financial support from Chic Group LTD, a manufacturer of electronic cigarettes in Poland. LK worked as an expert for the Polish National Committee for Standardization and for the European Committee for standardization of requirements and test methods for e-liquids and emissions. MLG receives fees for serving on an advisory board from Johnson & Johnson and grant support from Pfizer. BK declared no conflict of interest

## **Abstract**

Worldwide, cigarette smoking is a major cause of premature mortality and diseases that can be prevented. Given that people continue smoking despite the health risks, delivering nicotine without combustion should be considered a valuable and much less harmful way to reduce public health burden caused by smoking. E-cigarettes could play such a role if they are proved to be less harmful than combustible cigarettes. Although the number of clinical trials and human studies assessing safety of e-cigarettes is limited, numerous in vitro and in vivo studies reported the potential harmful effects of aerosol generated from e-cigarettes. This article reviews results of major clinical trials and laboratory studies with regard to cancer, cardiovascular and respiratory risk of e-cigarettes. Additionally, it also discusses the potential

application of e-cigarettes as smoking cessation tools. Most studies so far indicated that e-cigarettes are less harmful, but this applies only to smokers who completely switched to e-cigarettes. In the opinion of the authors, good quality research is important to establish the tolerance, safety, efficacy, and harm reduction potential of new technologies. Considering a significant role that physicians and other health providers play in helping smokers, there is an urgent need for evidence-based guidelines and recommendations for clinical practitioners on potential benefits and risks of e-cigarette use.

**Keywords:** e-cigarettes, tobacco harm reduction,

### **Recent outbreak of acute lung injuries related to vaping**

The year 2019 brought news about many tragic deaths linked to electronic cigarettes (e-cigarettes) use (vaping). By February 2020, the Centers for Disease Control and Prevention (CDC) reported 2758 hospitalizations for acute lung failure (EVALI; e-cigarette, or vaping, product use-associated lung injury), 64 of which ended in death [1,2]. The patients were mostly young people with a median age of 24 years who regularly used e-cigarettes. The first case was recorded in March 2019 with a peak number of hospital admissions reported in September 2019. After January 2020, the outbreak of the disease appears to diminish and only a few cases have been reported recently.

In view of these tragic events, the question of why this condition was not previously reported among regular e-cigarette users (common called “vapers”) has become relevant, despite the fact that vaping products have been available on the US market since 2007. The answer appeared quite quickly. The problem mainly occurred among e-cigarettes users who primarily vaped THC-containing products. Research showed that the e-cigarette refill fluids (e-liquids) used by hospitalized patients contained tetrahydrocannabinol (THC), a psychoactive substance derived from cannabis or cannabidiol (CBD), and were largely purchased on the black market.

The chest imaging of hospitalized EVALI patients showed, among others, acute eosinophilic pneumonia, diffuse alveolar damage, and lipid pneumonia [3]. Lipid pneumonia is an inflammatory reaction to the presence of lipids in the alveolar space, which usually results from inhalation of oily substances. Preliminary findings indicated that this oily substance was tocopherol acetate (vitamin E acetate), a lipophilic diluent used in some THC-containing e-liquids [4-6].

The evidence about causes of EVALI has become so prominent that on October 4, 2019 the United States Food and Drug Administration (FDA) issued a message which unambiguously warns to discontinue using vaping products containing THC and avoid purchasing e-cigarettes from unknown sources [7]. In addition, in December 2019 the United States Centers for Disease Control and Prevention (CDC) recommended refraining from adding any other substances (not intended by a manufacturer) to e-cigarette products, including those purchased through retail outlets [2].

### **What are E-cigarettes?**

Developed in 2003 by a Chinese pharmacist Hon Lik (Han Li), the electronic devices introducing nicotine to the user's body were, according to the inventor, meant to help quit smoking [8]. Although there is no combustion process involved, these new devices were named as electronic cigarettes or e-cigarettes mainly due to the first-generation products resembling a cigarette and the similar way of use. Despite some of the scientific community suggesting using the name Electronic Nicotine Delivery System (ENDS), the name e-cigarette has become widely accepted, which stigmatizes the device as another tobacco product. Since smokers are generally aware of the extreme harmfulness of tobacco products, e-cigarettes were also perceived as another potentially dangerous tobacco product. In addition, the limited availability of scientific publications after introduction of these products did not help potential users and health professionals to differentiate potential health risks of vaping compared to smoking.

Since the introduction of e-cigarettes to the global market, a significant number of studies that looked at potential health risk of vaping have been conducted and published. Currently, over 5300 records appear after typing the phrase "electronic cigarette" in the PubMed database. Many of those studies have looked at the relative risk of vaping compared to smoking. In general, studies suggest that use of e-cigarettes is associated with lower health risk compared to smoking tobacco cigarettes. Although this conclusion appears to be consistent across most of the studies, there are some limitations that need to be considered which we discuss below.

### **Limitations in the evaluation of e-cigarette safety**

One of the challenges in assessing the harmfulness of e-cigarettes are their rapidly changing design and product characteristics. For example, newer generation of e-cigarettes are significantly more effective in delivering nicotine to users than early generations of these products. Farsalinos et al. [9] demonstrated that the plasma nicotine concentration measured

after five minutes of smoking a conventional cigarette was 286% and 185% higher compared to the use of the first- and second-generation e-cigarettes, respectively. Currently, fourth generation e-cigarettes (Pod systems) may deliver nicotine to the same or even greater extent as conventional cigarettes, thus increasing the likelihood of addiction [10,11].

For some types of e-cigarettes the battery voltage can be adjusted—the higher the voltage, the higher the temperature of e-liquid. Substantial increase in vaporization temperature inside e-cigarette device may lead to thermal decomposition of two commonly used e-liquid solvents (glycerin and propylene glycol). Both solvents break down to formaldehyde, a toxic chemical with a proven carcinogenic potency. Kośmider et al. [12] demonstrated that an increase in the battery voltage from 3.0 V to 4.8 V causes more than tenfold increase in the amount of formaldehyde in the aerosol to a level similar to that in tobacco smoke. Another research showed that very low amounts of e-liquid on e-cigarette coil may result in overheating the device, and lead to a fifteen times higher emission of formaldehyde compared to traditional cigarette [13]. However, this work elicited a lot of controversy in the scientific community as the results presented were obtained during an unrealistic use of e-cigarette [14-17].

Another difficulty in assessing the potential risk associated with e-cigarettes use is the variety of flavoring substances present in e-liquids. In 2014 it was estimated that over 460 brands and over 7700 unique flavors of e-cigarettes were on the consumer market [18]. The authors of the discussed study also demonstrated that, unlike older brands emphasizing lesser harm compared to conventional cigarettes, manufacturers of newer e-cigarette brands focused more on various models and flavors options to choose from to increase their products appeal especially among young people. Such a great variety of product options results in safety concerns. For example, an aerosol generated from cherry-flavored liquids was found to contain more benzaldehyde than tobacco smoke. Benzaldehyde is a potential respiratory irritant that causes burning sensation in the nose and throat, coughing, and feeling of breathlessness [19].

E-liquids of sweet taste (e.g. toffee, milk, chocolate) may also contain prominent respiratory toxicants: diacetyl and 2,3-pentanedione [20]. Although these compounds are relatively safe when ingested, they have been shown to cause significant lung damage when inhaled. Inhalation of these compounds have been linked to reduced first-second intensive expiratory volume resulting in the development of bronchitis [21]. A condition called “Popcorn Lung” was reported among the employees of popcorn plants who were exposed to high concentration of diacetyl. Since some e-liquids contain the same toxicant, some authors suggested that the

use of e-cigarettes may also lead to the same disease [22]. However, the concentration of diacetyl in the traditional cigarette smoke is several hundred times higher and the “popcorn lung” disease has not been reported among regular smokers [23].

### **Toxic compound emissions from e-cigarettes and conventional cigarettes**

Tobacco smoke contains several thousand compounds, many of which have been shown to be toxic. On the other hand, aerosol generated from e-cigarettes contains a dozen to several dozen compounds. In 2012, the FDA published a list of 93 harmful and potentially harmful compounds (HPHCs) found in tobacco and tobacco smoke [24]. Out of these, only 5 compounds are present in the aerosol generated from the majority of commercially available e-liquids in quantities that may pose a potential health hazard to the user. These include acetaldehyde, acetone, acrolein, formaldehyde, and nicotine [25]. Lower yields of toxic compounds in the e-cigarette aerosol compared to tobacco smoke was also confirmed by a multicenter study, where authors found that the amount of toxic compounds present in the aerosol is 9 to 450 times lower than in tobacco smoke [26].

Margham et al. [27] compared the exposure to toxic compounds present in the aerosol of an e-cigarette and the standard 3R4F cigarette. This included the toxic compounds mentioned listed by the World Health Organization, FDA, and Health Canada. The reduction in the number and yields of toxic chemicals in aerosol compared to tobacco smoke was in the range of 92–99%. Also the US Surgeon General Report entirely devoted to e-cigarettes states that “E-cigarette aerosol is not harmless “water vapor,” although it generally contains fewer toxicants than combustible tobacco products.” At the same time, it adds that “the health effects and potentially harmful doses of heated and aerosolized constituents of e-cigarette liquids, including solvents, flavorants, and toxicants, are not completely understood” [28].

A significant reduction in toxic compound yields in e-cigarette aerosols compare to tobacco smoke as shown in the laboratory product testing, was confirmed in a population study with 5105 participants, including regular e-cigarette vapers and tobacco cigarette smokers. The urinary concentrations of biomarkers of exposure to toxic components of tobacco smoke, i.e., nicotine, tobacco-specific nitrosamines (NNK), metals, volatile organic compounds, and polycyclic aromatic hydrocarbons, were shown to be lower than those observed in current exclusive smokers and dual users of both products [29].

Potentially, the lower exposure to toxic substances from e-cigarette aerosols compared to tobacco smoke suggest that e-cigarettes are potentially less harmful products compared to

combustible cigarettes, which is the primary cause of premature mortality in developed countries mainly due to cancer, and cardiovascular and respiratory diseases [30]. Replacing combustible tobacco products with e-cigarettes would undoubtedly bring significant public health benefits if the reduction in toxicant exposure is proven to be associated with reduction in health risk.

### **E-cigarettes and a risk of cancer**

Smoking is the main cause of initiation of cancer process in the body resulting from several dozen carcinogenic compounds present in tobacco smoke, especially including 9 compounds classified by the International Agency of Research on Cancer (IARC) as group 1 carcinogens with proven carcinogenic effects on humans. Apart from traces of NNK found in e-cigarette liquids, there is one compound from this group, i.e., formaldehyde that has been reported in e-cigarette aerosols.

The carcinogenicity of various complex mixtures, including tobacco smoke and aerosol emitted from e-cigarettes, can only be established on the basis of a documented relationship between the exposure to the factor and increase in the incidence of cancer in exposed humans or animals (epidemiological studies). Such a relationship was documented for smokers but not for e-cigarette users, as these products are introduced recently on the consumer market. The reason behind this is the long cancer latency periods ranging from several to several dozen years. Although no epidemiological studies are available at the moment, we can roughly assess the relative cancer risk of e-cigarette by measuring biomarkers of exposure to carcinogens and compare the measured values to exposure levels observed in smokers and non-smokers.

In one of the first exposure assessment studies, traditional smokers completely substituted their tobacco cigarettes form e-cigarettes for two weeks. That study showed a drastic reduction in exposure to carcinogens, including 1,3-butadiene, benzene, acrylonitrile, and NNK. Concentrations of the corresponding biomarkers decreased on average by 57% after the first week and 67% after the second week [31].

A cross-sectional study of a group of 181 volunteers showed that former cigarette smokers who gave up smoking in favor of e-cigarettes for at least six months had significantly reduced exposure to carcinogenic and toxic compounds compared to those who continued smoking. Levels of biomarkers of exposure measured in e-cigarette users were similar to those subjects

who used nicotine replacement therapy (NRT) [32]. Importantly, no reduction in exposure was observed in persons who concurrently smoked cigarettes and vaped e-cigarettes.

Cadmium is one of the IARC group 1 carcinogens present in the tobacco smoke. A cross-sectional study of 156 volunteers showed that after switching from cigarettes to e-cigarettes, a 69% decrease was observed in the concentration of cadmium in blood after six months [33].

The most comprehensive, and probably the most representative up-to-date, report of the US National Academies of Sciences, Engineering, and Medicine (NASEM) states that the current evidence related to the potential link between e-cigarette use and cancer is too scarce to allow meaningful conclusions to be drawn about the cancer process or about intermediate clinical endpoints [34].

### **E-cigarettes and a risk of Cardiovascular diseases**

According to the US Surgeon General Report, the mechanism through which smoking affects the cardiovascular system is complex but well documented [35]. It was proven that nicotine, carbon monoxide, and tobacco combustion products are primarily responsible for cardiovascular dysfunction in smokers.

Nicotine activates the sympathetic system and releases catecholamines, adrenaline, and noradrenaline, which increase the myocardial function and oxygen demand by rapidly increasing blood pressure, accelerating heart rate, and vasospasm. There is no reason to believe that nicotine in the e-cigarette aerosols affects the cardiovascular system differently than nicotine in tobacco smoke.

Carbon monoxide reacts with hemoglobin to form carboxyhemoglobin, which is not capable of carrying oxygen, resulting in a reduction in oxygen availability. However, this does not happen with e-cigarettes because of the absence of combustion.

Combustion products of tobacco, such as polycyclic aromatic hydrocarbons (PAHs), oxidizing compounds, free radicals, and respirable particles, play a significant role in this process. According to the HPHCs list, toxic effects on the cardiovascular system can be attributed to at least 12 compounds present in tobacco smoke. Only two of these, acrolein and propionic aldehyde, are present in e-cigarette aerosol due to thermal decomposition of the solvents. However, some cardiovascular toxicants that are not present in e-liquids could be generated from flavorings and additives during vaporization. For example, benzene has been shown to be generated from certain flavorings used in e-liquids when e-cigarette users takes puffs on the device [36].



To date, the comparison of solid particles in tobacco smoke and aerosol remains controversial. Their composition and characteristics in aerosol differ significantly from those in tobacco smoke. At the same time, the data to determine their cardiovascular toxicity is insufficient [37].

Combustion products induce inflammation which activates platelets and leads to vascular endothelial dysfunction. Many researchers believe that endothelial dysfunction is an essential element to the future development of atherosclerotic damage. Hence it is designated as “The risk of the risk factors” [38]. By quitting tobacco products and switching to e-cigarettes, the smoker greatly reduces the inhalation of compounds resulting from tobacco combustion. Consequently, the negative effect of the aerosol on vascular endothelial dysfunction is expected to be reduced. This has been confirmed by an extensive review by Knura et al. [39]. Based on the available data, the authors concluded that e-cigarette aerosol damages the vascular endothelium, but to a lesser extent than tobacco smoke.

Using an e-cigarette also causes oxidative stress due to oxidative compounds but comparatively lesser than a conventional cigarette [40].

Randomized clinical study, perceived as the “gold standard” for analysis of treatment results by many, showed that flow-induced vasodilatation in the brachial artery significantly increases one month after switching to e-cigarettes [41].

Based on the previous clinical studies, Benowitz et al. [42,43] concluded that the overall acute circulatory failure associated with e-cigarettes is consistent with the nicotine effect. Additionally, the authors believe that the cardiovascular risk caused by nicotine inhaled from e-cigarettes is quite low in people without preexisting cardiovascular disease. However, people with diagnosed cardiovascular disease are still prone to the risk but comparatively lower than the traditional smokers. If conventional cigarettes are completely replaced by e-cigarettes, the harmfulness of smoking can be significantly reduced and smokers who completely switch to e-cigarettes would experience reduction in cardiovascular risk.

According to the NASEM report, there is no evidence available to confirm whether the use of e-cigarettes is associated with ischemic heart disease, stroke and peripheral artery disease, and subclinical arteriosclerosis. There is also insufficient evidence that the use of e-cigarettes is associated with long-term changes in heart rate, blood pressure, and heart geometry and function. However, there is convincing evidence that heart rate increases shortly after nicotine

inhalation from e-cigarettes. There is also moderate evidence that diastolic blood pressure increases shortly after inhaling nicotine from e-cigarettes [34].

Although current, but still limited, literature suggests that the use of e-cigarettes may lead to less negative cardiovascular effects than traditional cigarettes, some researchers see a need for additional, high quality randomized controlled trials to clearly establish the cardiovascular safety of e-cigarettes. Future studies should continue to focus on the study of both long- and short-term effects of exposure to e-cigarettes and their potential role in the development of cardiovascular disease [44].

### **E-cigarettes and a risk of Respiratory diseases**

As e-cigarettes are products purported for inhalation use, one may expect that any potential risk of these products would be reflected particularly in respiratory dysfunction. The e-cigarette users inhale relatively large amounts of two substances that are only present in small quantities in tobacco smoke. These substances are glycerin and propylene glycol that served as nicotine solvents and carriers in e-cigarettes and as humectants in tobacco cigarettes. In e-liquids they constitute about 80% of the overall content of e-liquids. Although these compounds are commonly used in cosmetics, pharmaceutical, and food industries, the long-term inhalation risk has not been well studied previously.

The longest study (3.5 years) looked at changes in respiratory symptoms in e-cigarettes users and non-users. Changes in hemodynamic parameters, respiratory functions, and high-resolution computed tomography of lungs were evaluated in 31 volunteers who never smoked cigarettes but started and continued to use e-cigarettes. No significant changes were observed compared to the control group of persons who never smoked [45].

A two-year study of a group of 209 e-cigarettes users included examination of vital parameters, electrocardiography, lung-function testing, exposure to nicotine and selected compounds, and urges to smoke a cigarette and withdrawal effects. No serious health events were observed. The most common negative health symptoms reported by the volunteers were headache (28.7%), nasopharyngeal inflammation (19.6%), sore throat, and cough (16.9%) [46].

A systematic review of clinical data related to the effects of e-cigarettes use on the respiratory system found that replacing combustible tobacco cigarettes with e-cigarettes significantly

improved health outcomes in patients with chronic obstructive pulmonary disease (COPD) and chronic asthma [47].

In contrast, a representative national (US) longitudinal study from 2013–2016 found that e-cigarettes appear to be an independent risk factor for respiratory diseases. Among those who did not report respiratory diseases (COPD, chronic bronchitis, emphysema or asthma), longitudinal analysis revealed statistically significant relationships between previous (AOR = 1.31) and current use of e-cigarettes (AOR = 1.29) and lung diseases. Smoking was also significantly associated with any respiratory disease but the odds was twice as high as for vaping (AOR = 2.56). An even higher value was recorded for dual users (AOR = 3.30). The authors concluded that switching from combustible tobacco products, including cigarettes, to e-cigarettes may reduce the risk of developing respiratory diseases. At the same time, the authors pointed to a high incidence among smokers using both products, which is associated with an increased risk beyond the use of combustible tobacco products. This is a prevailing argument that dual use of e-cigarettes and tobacco cigarettes does not reduce a respiratory risk from smoking but may actually increase risk such risk. [48].

Similar conclusions are included in the NASEM report, particularly with regard to dual users. Although the report concluded that there is no evidence available that e-cigarettes cause respiratory diseases in humans, it also states that there is limited evidence of improved lung function and respiratory symptoms in adult asthma smokers who switch to e-cigarettes in full or in part (dual use). This also applies to adult smokers with obstructive lung disease [34].

### **E-cigarettes as a cigarette smoking cessation tool**

Although the intension of e-cigarette inventor, Chinese pharmacist Hon Lik, was to develop a device for smokers that would help them quit smoking, only few clinical trials and population-based studies evaluated efficacy of e-cigarettes for smoking cessation. The potential effectiveness of e-cigarettes in helping smokers to quit smoking is an area of much controversy. This is perfectly illustrated by two meta-analyses published by Kalkhoran & Glantz [49] and Malas et al. [50].

The first team analyzed 30 studies and the second 38 studies, 25 of which were analyzed by both research teams. From these 25 studies, the first team excluded 10 works due to the lack of a control group of people who did not use e-cigarettes. The results obtained were diametrically different. In the first study, the odds ratio was OR = 0.72 (95% CI; 0.57– 0.91), leading authors to the conclusion that use of e-cigarettes among smokers reduces their

chances of quitting. In the second meta-analysis, the odds of quitting were higher (OR = 1.63 (95% CI; 1.17–2.27)) suggesting that e-cigarette users are more likely to quit smoking than smoker who does not use these products. This is an example of how different methodological approaches can lead to contradicting conclusions.

NASEM report included an analysis of studies and clinical trials published until the beginning of 2018 [34]. The authors conclude that there is limited evidence that e-cigarettes can be effective in promoting and supporting smoking cessation and moderate evidence from randomized clinical trials that e-cigarettes with nicotine are more effective than nicotine-free e-cigarettes. Observational studies showed that more frequent use of e-cigarettes is associated with an increased likelihood of quitting smoking. However, there is insufficient evidence from randomized studies on the efficacy of e-cigarettes as quit-smoking aids compared to the absence of anti-nicotine therapies or FDA-approved treatments.

The conclusions cited above about limited efficacy of e-cigarettes as a tool to help smokers quit cigarette use need to be updated since two comprehensive and large randomized clinical trials have been published recently [51,52]. In the first trial, 886 participants were randomized to e-cigarettes or nicotine replacement therapy (NRTs). Both groups were provided with a supportive behavioral therapy. The 1-year abstinence rate was 18.0% in the e-cigarette group compared to 9.9% in the NRT group. The second trial involved 1124 volunteers and compared efficacy of e-cigarettes combined with nicotine patches compared to nicotine patches alone. Six-month abstinence was 7% versus 2%, respectively. Given that the pharmacokinetic profiles of nicotine from smoking and vaping are similar, it could be assumed that combination therapy (e-cigarette plus NRT) can become an important strategy to improve smoking cessation rates.

A recently released report by the US Surgeon General [53] on the smoking cessation also evaluated potential application of e-cigarettes as smoking-cessation tools. The conclusions were consistent with those of the NASEM report cited above. The authors of the report pointed out that e-cigarettes can be attractive to traditional cigarette smokers because their use reflects some of the sensorimotor characteristics of traditional cigarette smoking, including stimulation of the upper respiratory tract, sensations, and taste of the e-cigarette aerosol in the mouth, hand movements, inhalation and exhalation of the aerosol that visually often resembles cigarette smoke. Given the potentially important role of such factors in the abuse liability of tobacco cigarettes, their presence in the process of vaping may actually make e-

cigarettes more attractive to smokers who may see those products as better substitutes for cigarettes than NRT.

It becomes particularly important to assess the scientific evidence on the impact of e-cigarettes on smoking cessation among adult smokers in the context of significant uptake of e-cigarettes by youths. The popularity of e-cigarettes among adolescents and young adults has increased drastically in recent years, particularly after the introduction of the fourth generation of e-cigarettes in the shape of USB flash drives [54].

### **Summary**

The skyrocketing use of e-cigarettes among youth and recent EVALI epidemic observed within the last year have intensified the controversy around e-cigarettes and questioned their potential for public health benefits as a substitute for traditional cigarettes. Numerous opinions and editorial pieces as it relates to public health consequences of e-cigarette use have been published by many prestigious and influential medical journals.

The Lancet's editorial stated that there is no firm evidence to claim that e-cigarettes are healthier than cigarettes or can support quitting smoking. In addition, the renormalization of smoking in the form of e-cigarettes, not only among smokers but also among young people and nonsmokers, poses a risk to the entire population from nicotine use and may lead to large-scale addiction [55]. This article was criticized by Professor JN Newton, Director of Health Improvement at Public Health England, who stated that there is an international consensus that the use of e-cigarettes is likely to be much less harmful than smoking and that e-cigarettes have an important role to play in tobacco control [56].

Beaglehole et al. [57] in their commentary published also by The Lancet considered e-cigarettes as a strategy to minimizing harm caused by smoking. E-cigarettes have a potential to be used as a complementary harm reduction strategy, but currently this strategy is underestimated because achieving total nicotine abstinence is the overriding goal for many regulators and public health advocates. In the US the effectiveness to achieve total nicotine abstinence in years 2014-2015 was significantly higher than that for 2010-2011 (5.6 vs 4.5). Taking into account, that e-cigarettes started to gain popularity about 2010, increase in total nicotine abstinence might be explained by increasing their use in the population. This is supported by the fact, that 38.2% of current smokers and 49.3% of recent quitters had tried e-cigarettes, and 11.5% and 19.0% used them currently (every day or some days) [58]. Those data suggest, that e-cigarettes have a potential as harm reduction product, what should be

considered especially as currently used methods for nicotine addiction treatment used by providers have limited effectiveness and result in low quit rates.

The authors of the article in *Science* expressed opinions that regulatory approaches to e-cigarettes in the absence of scientific evidence certainty must include many compromises. Finding the optimal balance between providing smokers with fully regulated nicotine vaping products, while at the same time significantly reducing the risk of using e-cigarettes in young people appears to be a proper compromise. In this context, the authors cite the example of the UK, which adopted nicotine vaping harm reduction as a safer alternative to combustible products, and was able to make appropriate regulations that led to a consensus between the risks to young people and the benefits to smokers [59].

The authors of the article in *JAMA* emphasize that health care providers should remember that smoking remains a major public health threat. They point out the significant reduction in the prevalence of smoking after introduction of e-cigarettes. Indeed, the smoking prevalence has reached its lowest level in many countries despite a sharp increase in the use of e-cigarettes, and perhaps partially because of this. It is important that former cigarette smokers who use e-cigarettes do not return to traditional cigarettes, which are likely to present a much higher risk than e-cigarettes [60].

There are a number of questions about e-cigarettes that lack answers: whether use of e-cigarettes among youth will lead to smoking (so called “gateway effect”), whether banning flavored e-cigarettes will discourage young people from using them, whether a reducing nicotine levels in e-cigarettes will encourage smokers who have completely switched to e-cigarettes to return to combustible tobacco products, to what extent long-term vaping affects the health of the user.

In our view, good quality clinical trials will be increasingly important to establish safety, efficacy, and harm reduction potential for new technologies. Despite many areas of ambiguity, current evidence suggest that e-cigarettes are less harmful than combustible products, but this only applies to smokers who completely switched to e-cigarettes. Thus, e-cigarettes still hold a great potential to reduce incidences of tobacco-related diseases and could be a part of the strategy to reduce the damage caused by smoking. Therefore, mechanisms should be developed to protect young people from using e-cigarettes but support smokers in their decisions to quit smoking with e-cigarettes. Considering the importance of this topic for public health, we encourage internists and general practitioners to expand their

knowledge about e-cigarettes as new evidence on the impact of e-cigarettes on users' health will emerge in future.

#### Acknowledgments:

This work was supported by Medical University of Silesia in Katowice (Poland), Contract No. KNW-1-039/K/9/O (AS, LK) and was supported by grant from the US National Cancer Institute and the US Center for Tobacco Products of the Food and Drug Administration (U54CA228110) (MLG). The views and opinions expressed in this letter are those of the authors and do not necessarily represent the official position of the National Institutes of Health or the Food and Drug Administration.

#### References

1. Outbreak of Lung Injury Associated with the Use of E-Cigarette, or Vaping, Products. Centers for Disease Control and Prevention (CDC).  
[https://www.cdc.gov/tobacco/basic\\_information/e-cigarettes/severe-lung-disease.html](https://www.cdc.gov/tobacco/basic_information/e-cigarettes/severe-lung-disease.html)  
[online access 2.02.2020](#)
2. Krishnasamy VP, Hallowell BD, Ko JY, et al. Update: Characteristics of a Nationwide Outbreak of E-cigarette, or Vaping, Product Use-Associated Lung Injury - United States, August 2019-January 2020. *MMWR Morb Mortal Wkly Rep.* 2020; 69: 90-94.
3. Henry TS, Kanne JP, Kligerman SJ. Imaging of Vaping-Associated Lung Disease. *N Engl J Med* 2019; 381: 1486-1487.
4. Blount BC, Karwowski MP, Morel-Espinosa M, et al. Evaluation of bronchoalveolar lavage fluid from patients in an outbreak of e-cigarette, or vaping, product use-associated lung injury — 10 states, August–October 2019. *MMWR Morb Mortal Wkly Rep.* 2019; 68: 1040-1041.
5. Blount BC, Karwowski MP, Shields PG, et al. Vitamin E acetate in bronchoalveolar-lavage fluid associated with EVALI. *N Engl J Med.* 2020; 382: 697-705.
6. Bhat TA, Kalathil SG, Bogner PN, et al. An Animal Model of Inhaled Vitamin E Acetate and EVALI-like Lung Injury. *N Engl J Med.* DOI: 10.1056/NEJMc2000231
7. Vaping Illness Update: FDA Warns Public to Stop Using Tetrahydrocannabinol (THC)-Containing Vaping Products and Any Vaping Products Obtained Off the Street. US Food & Drug Administration <https://www.fda.gov/consumers/consumer->

[updates/vaping-illness-update-fda-warns-public-stop-using-tetrahydrocannabinol-the-containing-vaping](#)

8. Li, H. Non-Combustible Electronic Atomizing Cigarette. CN Patent No. 2643681, 2004.
9. Farsalinos KE, Spyrou A, Tsimopoulou K, et al. Nicotine absorption from electronic cigarette use: comparison between first and new-generation devices. *Sci Rep.* 2014; 4: 4133.
10. Talih S, Salman R, El-Hage R, et al. Characteristics and toxicant emissions of JUUL electronic cigarettes. *Tob Control.* 2019; 28: 678-680.
11. Goniewicz ML, Boykan R, Messina CR, et al. High exposure to nicotine among adolescents who use Juul and other vape pod systems ('pods'). *Tob Control.* 2019; 28: 676-677.
12. Kosmider L, Sobczak A, Fik M, et al. Carbonyl compounds in electronic cigarette vapors: effects of nicotine solvent and battery output voltage. *Nicotine Tob Res.* 2014; 16: 1319-1326.
13. Jensen RP, Luo W, Pankow JF, et al. Hidden formaldehyde in e-cigarette aerosols. *N Engl J Med.* 2015; 372: 392-394.
14. Farsalinos KE, Voudris V, Spyrou A, Poulas K. E-cigarettes emit very high formaldehyde levels only in conditions that are aversive to users: A replication study under verified realistic use conditions. *Food Chem Toxicol.* 2017; 109: 90-94.
15. Bates CD, Farsalinos KE. E-cigarettes need to be tested for safety under realistic conditions. *Addiction.* 2015; 110: 1688-1689.
16. Kershaw O. 'Dry puff' and electronic cigarettes. *Addiction.* 2015; 110(12): 2038.
17. Nitzkin JL, Farsalinos K, Siegel M. More on hidden formaldehyde in e-cigarette aerosols. *N Engl J Med.* 2015; 372(16): 1575.
18. Zhu SH, Sun JY, Bonnevie E, et al. Four hundred and sixty brands of e-cigarettes and counting: implications for product regulation. *Tob Control.* 2014; 23 Suppl 3:iii3-9.
19. Kosmider L, Sobczak A, Prokopowicz A, et al. Cherry-flavoured electronic cigarettes expose users to the inhalation irritant, benzaldehyde. *Thorax.* 2016; 71: 376-377.
20. Farsalinos KE, Kistler KA, Gillman G, Voudris V. Evaluation of electronic cigarette liquids and aerosol for the presence of selected inhalation toxins. *Nicotine Tob Res.* 2015; 17: 168-174.



21. van Rooy FG1, Rooyackers JM, Prokop M, et al. Bronchiolitis obliterans syndrome in chemical workers producing diacetyl for food flavorings. *Am J Respir Crit Care Med.* 2007;176: 498-504.
22. Chung S, Baumlin N, Dennis JS, et al. Electronic Cigarette Vapor with Nicotine Causes Airway Mucociliary Dysfunction Preferentially via TRPA1 Receptors. *Am J Respir Crit Care Med.* 2019; 200: 1134-1145.
23. Fujioka K1, Shibamoto T. Determination of toxic carbonyl compounds in cigarette smoke. *Environ Toxicol.* 2006; 21: 47-54.
24. Food and Drug Administration. Harmful and potentially Harmful Constituents in Tobacco Products and Tobacco Smoke; Established List. *Federal Register.* 2012; 77 (64): 20034-7
25. Tayyarah R, Long GA. Comparison of select analytes in aerosol from e-cigarettes with smoke from conventional cigarettes and with ambient air. *Regul Toxicol Pharmacol.* 2014; 70: 704-710.
26. Goniewicz ML, Knysak J, Gawron M, et al. Levels of selected carcinogens and toxicants in vapour from electronic cigarettes. *Tob Control.* 2014; 23:133-139.
27. Margham J, McAdam K, Forster M, et al. Chemical Composition of Aerosol from an E-Cigarette: A Quantitative Comparison with Cigarette Smoke. *Chem Res Toxicol.* 2016; 29: 1662-1678.
28. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health (2016). E-Cigarette Use Among Youth and Young Adults. A Report of the Surgeon General. 2016: p. 125.
29. Goniewicz ML, Smith DM, Edwards KC, et al. Comparison of Nicotine and Toxicant Exposure in Users of Electronic Cigarettes and Combustible Cigarettes. *JAMA Netw Open.* 2018; 1: e185937
30. World Health Organization (WHO): Global health risks: mortality and burden of disease attributable to selected major risks. Geneva, 2009 ISBN 978 92 4 156387 1
31. Goniewicz ML, Gawron M, Smith DM, et al. Exposure to Nicotine and Selected Toxicants in Cigarette Smokers Who Switched to Electronic Cigarettes: A Longitudinal Within-Subjects Observational Study. *Nicotine Tob Res.* 2017; 19: 160-167.

32. Shahab L, Goniewicz ML, Blount BC, et al. Nicotine, Carcinogen, and Toxin Exposure in Long-Term E-Cigarette and Nicotine Replacement Therapy Users: A Cross-sectional Study. *Ann Intern Med.* 2017; 166: 390-340.
33. Prokopowicz A, Sobczak A, Szula-Chraplewska M, et al. Exposure to Cadmium and Lead in Cigarette Smokers Who Switched to Electronic Cigarettes. *Nicotine Tob Res.* 2019; 21: 1198-1205.
34. National Academies of Sciences, Engineering, and Medicine. 2018. Public health consequences of e-cigarettes. Washington, DC: The National Academies Press. doi:<https://doi.org/10.17226/24952>.
35. U.S. Department of Health and Human Services. The Health Consequences of Smoking: 50 Years of Progress. A Report of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2014.
36. Pankow JF, Kim K, McWhirter KJ, et al. Benzene formation in electronic cigarettes. *PLoS One.* 2017; 12: e0173055.
37. Floyd EL, Queimado L, Wang J, et al. Electronic cigarette power affects count concentration and particle size distribution of vaping aerosol. *PLoS One.* 2018; 13: e0210147.
38. Bonetti PO, Lerman LO, Lerman A. Endothelial dysfunction. a marker of atherosclerotic risk. *Arterioscler Thromb Vasc Biol.* 2003; 23: 168-175.
39. Knura M, Dragon J, Łabuzek K, Okopień B. [The impact of electronic cigarettes usage on the endothelial function and the progression of atherosclerosis]. *Pol Merkur Lekarski.* 2018; 44: 26-30. (Polish)
40. Ikonomidis I, Vlastos D, Kourea K, et al. Electronic Cigarette Smoking Increases Arterial Stiffness and Oxidative Stress to a Lesser Extent Than a Single Conventional Cigarette: An Acute and Chronic Study. *Circulation.* 2018; 137: 303-306.
41. George J, Hussain M, Vadiveloo T, et al. Cardiovascular Effects of Switching From Tobacco Cigarettes to Electronic Cigarettes. *J Am Coll Cardiol.* 2019; 74: 3112-3120.
42. Benowitz NL, Burbank AD. Cardiovascular toxicity of nicotine: Implications for electronic cigarette use. *Trends Cardiovasc Med.* 2016; 26: 515-523.
43. Benowitz NL, Fraiman JB. Cardiovascular effects of electronic cigarettes. *Nat Rev Cardiol.* 2017; 14: 447-456.

44. Buchanan ND, Grimmer JA, Tanwar V, et al. Cardiovascular risk of electronic cigarettes: a review of preclinical and clinical studies. *Cardiovasc Res.* 2020; 116: 40-50.
45. Polosa R, Cibella F, Caponnetto P, et al. Health impact of E-cigarettes: a prospective 3.5-year study of regular daily users who have never smoked. *Sci Rep.* 2017; 7: 13825.
46. Walele T, Bush J, Koch A, et al. Evaluation of the safety profile of an electronic vapour product used for two years by smokers in a real-life setting. *Regul Toxicol Pharmacol.* 2017; 92: 226-238.
47. Morjaria JB, Mondati E, Polosa R. E-cigarettes in patients with COPD: current perspectives. *Int J Chron Obstruct Pulmon Dis.* 2017; 12: 3203-3210.
48. Bhatta DN, Glantz SA. Association of E-Cigarette Use With Respiratory Disease Among Adults: A Longitudinal Analysis. *Am J Prev Med.* 2020; 58: 182-190.
49. Kalkhoran S, Glantz SA. E-cigarettes and smoking cessation in real-world and clinical settings: a systematic review and meta-analysis. *Lancet Respir Med.* 2016; 4: 116-128.
50. Malas M, van der Tempel J, Schwartz R, et al. Electronic Cigarettes for Smoking Cessation: A Systematic Review. *Nicotine Tob. Res.* 2016; 18: 1926-1936.
51. Hajek P, Phillips-Waller A, Przulj D, et al. A Randomized Trial of E-Cigarettes versus Nicotine-Replacement Therapy. *N Engl J Med.* 2019; 380: 629–637.
52. Walker N, Parag V, Verbiest M, et al. Nicotine patches used in combination with e-cigarettes (with and without nicotine) for smoking cessation: a pragmatic, randomised trial. *Lancet Respir Med.* 2020; 8: 54–64.
53. U.S. Department of Health and Human Services. Smoking Cessation. A Report of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2020.
54. Cullen KA, Gentzke AS, Sawdey MD, et al. e-Cigarette Use Among Youth in the United States, 2019. *JAMA.* 2019. doi: 10.1001/jama.2019.18387.
55. The Lancet. E-cigarettes: time to realign our approach? *Lancet* 2019; 394: 1297.
56. Newton JN. Time for The Lancet to realign with the evidence on e-cigarettes? *Lancet.* 2019; 394: 1804-1805.
57. Beaglehole R, Bates C, Youdan B, Bonita R. Nicotine without smoke: fighting the tobacco epidemic with harm reduction. *Lancet.* 2019; 394: 718-720.

58. Zhu SH, Zhuang YL, Wong S, et al. E-cigarette use and associated changes in population smoking cessation: evidence from US current population surveys. *BMJ*. 2017; 358: j3262.
59. Fairchild A, Healton C, Curran J, et al. Evidence, alarm, and the debate over e-cigarettes. *Science*. 2019; 366: 1318-1320.
60. Baldassarri SR, Fiellin DA, Friedman AS. Vaping-Seeking Clarity in a Time of Uncertainty. *JAMA*. 2019. doi: 10.1001/jama.2019.16493.